arXiv:1607.03503v1 [math.GR] 12 Jul 2016

A BIEBERBACH THEOREM FOR CRYSTALLOGRAPHIC GROUP EXTENSIONS

JOHN G. RATCLIFFE AND STEVEN T. TSCHANTZ

Abstract. In this paper we prove that for each dimension n there are only finitely many isomorphism classes of pairs of groups (Γ, N) such that Γ is an n-dimensional crystallographic group and N is a normal subgroup of Γ such that Γ/N is a crystallographic group.

1. Introduction

An *n-dimensional crystallographic group* (*n-space group*) is a discrete group Γ of isometries of Euclidean n-space E^n whose orbit space E^n/Γ is compact. The 3-space groups are the symmetry groups of crystalline structures, and so are of fundamental importance in the science of crystallography.

In response to Hilbert's Problem 18, L. Bieberbach [2] proved that for each dimension n there are only finitely many isomorphism classes of n-space groups. In this paper we prove a relative version of Bieberbach's theorem. We prove that for each dimension n there are only finitely many isomorphism classes of pairs of groups (Γ, N) such that Γ is an n-space group and N is a normal subgroup of Γ such that Γ/N is a space group.

Our relative Bieberbach theorem has a geometric interpretation in the theory of flat orbifolds. By Theorems 7, 8, and 10 of [8] the isomorphism classes of pairs of groups (Γ, N) such that Γ is an n-space group and N is a normal subgroup of Γ such that Γ/N is a space group correspond to the affine equivalence classes of geometric orbifold fibrations of compact, connected, flat n-orbifolds. Therefore, our relative Bieberbach theorem is equivalent to the theorem that for each dimension n there are only finitely many affine equivalence classes of geometric orbifold fibrations of compact, connected, flat n-orbifolds. This is known for $n = 3$ by the work of Conway-Friedrichs-Huson-Thurston[4] and Ratcliffe-Tschantz [8].

We now outline the proof of our relative Bieberbach theorem. Let m be a positive integer less than n. Let M be an m-space group and let Δ be an $(n-m)$-space group. Let $\mathrm{Iso}(\Delta, \mathrm{M})$ be the set of isomorphism classes of pairs (Γ, N) where N is a normal subgroup of an n-space group Γ such that N is isomorphic to M and Γ/N is isomorphic to Δ. As there are only finitely many isomorphism classes of the groups Δ and M by Bieberbach's theorem [2], it suffices to prove that $\mathrm{Iso}(\Delta, \mathrm{M})$ is finite.

Next, we define a set $\mathrm{Out}(\Delta, \mathrm{M})$ in terms of $\mathrm{Out}(\Delta)$ and $\mathrm{Out}(\mathrm{M})$. That the set $\mathrm{Out}(\Delta, \mathrm{M})$ is finite follows easily from a theorem of Baues and Grunewald [1] that the outer automorphism group of a crystallographic group is an arithmetic group. We define a function $\omega : \mathrm{Iso}(\Delta, \mathrm{M}) \to \mathrm{Out}(\Delta, \mathrm{M})$. We prove that $\mathrm{Iso}(\Delta, \mathrm{M})$ is finite by showing that the fibers of ω are finite by a cohomology of groups argument.

2. Normal Subgroups of Space Groups

A map $\phi : E^n \to E^n$ is an isometry of E^n if and only if there is an $a \in E^n$ and an $A \in \mathrm{O}(n)$ such that $\phi(x) = a + Ax$ for each x in E^n. We shall write $\phi = a + A$. In particular, every translation $\tau = a + I$ is an isometry of E^n.

Let Γ be an n-space group. Define $\eta : \Gamma \to \mathrm{O}(n)$ by $\eta(a + A) = A$. Then η is a homomorphism whose kernel is the group T of translations in Γ. The image of η is a finite group Π called the *point group* of Γ.

Let H be a subgroup of an n-space group Γ. Define the *span* of H by the formula

$$\mathrm{Span(H)} = \mathrm{Span}\{a \in E^n : a + I \in \mathrm{H}\}.$$

Note that Span(H) is a vector subspace V of E^n. Let $V^\perp$ denote the orthogonal complement of V in E^n.

Theorem 1. (Theorem 2 [8]) *Let* N *be a normal subgroup of an n-space group Γ, and let $V = \mathrm{Span(N)}$.*

1. *If $b + B \in \Gamma$, then $BV = V$.*
2. *If $a + A \in \mathrm{N}$, then $a \in V$ and $V^\perp \subseteq \mathrm{Fix}(A)$.*
3. *The group* N *acts effectively on each coset $V + x$ of V in E^n as a space group of isometries of $V + x$.*

Let Γ be an n-space group. The *dimension* of Γ is n. If N is a normal subgroup of Γ, then N is a m-space group with $m = \dim(\mathrm{Span(N)})$ by Theorem 1(3).

Definition: Let N be a normal subgroup N of an n-space group Γ, and let $V = \mathrm{Span(N)}$. Then N is said to be a *complete normal subgroup* of Γ if

$$\mathrm{N} = \{a + A \in \Gamma : a \in V \text{ and } V^\perp \subseteq \mathrm{Fix}(A)\}.$$

Lemma 1. (Lemma 1 [8]) *Let* N *be a complete normal subgroup of an n-space group Γ, and let $V = \mathrm{Span(N)}$. Then Γ/N acts effectively as a space group of isometries of E^n/V by the formula* $(\mathrm{N}(b + B))(V + x) = V + b + Bx$.

Remark 1. A normal subgroup N of a space group Γ is complete precisely when Γ/N is a space group by Theorem 5 of [8].

Let N be a complete normal subgroup of an n-space group Γ, let $V = \mathrm{Span(N)}$, and let $V^\perp$ be the orthogonal complement of V in E^n. Let $\gamma \in \Gamma$. Then $\gamma = b + B$ with $b \in E^n$ and $B \in \mathrm{O}(n)$. Write $b = \overline{b} + b'$ with $\overline{b} \in V$ and $b' \in V^\perp$. Let $\overline{B}$ and B' be the orthogonal transformations of V and $V^\perp$, respectively, obtained by restricting B. Let $\overline{\gamma} = \overline{b} + \overline{B}$ and $\gamma' = b' + B'$. Then $\overline{\gamma}$ and γ' are isometries of V and $V^\perp$, respectively.

Euclidean n-space E^n decomposes as the Cartesian product $E^n = V \times V^\perp$. Let $x \in E^n$. Write $x = v + w$ with $v \in V$ and $w \in V^\perp$. Then

$$(b + B)x = b + Bx = \overline{b} + b' + Bv + Bw = (\overline{b} + \overline{B}v) + (b' + B'w).$$

Hence the action of Γ on E^n corresponds to the diagonal action of Γ on $V \times V^\perp$ defined by the formula

$$\gamma(v, w) = (\overline{\gamma}v, \gamma' w).$$

Here Γ acts on both V and $V^\perp$ via isometries. The kernel of the corresponding homomorphism from Γ to $\mathrm{Isom}(V)$ is the group

$$\mathrm{K} = \{b + B \in \Gamma : b \in V^\perp \text{ and } V \subseteq \mathrm{Fix}(B)\}.$$

We call K the *kernel of the action* of Γ on V. The group K is a normal subgroup of Γ. The action of Γ on V induces an effective action of Γ/K on V via isometries. The group Γ/K acts on V as a discrete group of isometries if and only if Γ/NK is a finite group by Theorem 3(4) of [10].

The group N is the kernel of the action of Γ on $V^\perp$, and so the action of Γ on $V^\perp$ induces an effective action of Γ/N on $V^\perp$ via isometries. Orthogonal projection from E^n to $V^\perp$ induces an isometry from E^n/V to $V^\perp$. Hence Γ/N acts on $V^\perp$ as a space group of isometries by Lemma 1.

Let $\overline{\Gamma} = \{\overline{\gamma} : \gamma \in \Gamma\}$. If $\gamma \in \Gamma$, then $(\overline{\gamma})^{-1} = \overline{\gamma^{-1}}$, and if $\gamma_1, \gamma_2 \in \Gamma$, then $\overline{\gamma_1}\,\overline{\gamma_2} = \overline{\gamma_1\gamma_2}$. Hence $\overline{\Gamma}$ is a subgroup of $\mathrm{Isom}(V)$. The map $\mathrm{B} : \Gamma \to \overline{\Gamma}$ defined by $\mathrm{B}(\gamma) = \overline{\gamma}$ is an epimorphism with kernel K. The group $\overline{\Gamma}$ is a discrete subgroup of $\mathrm{Isom}(V)$ if and only if Γ/NK is finite by Theorem 3(4) of [10].

Let $\Gamma' = \{\gamma' : \gamma \in \Gamma\}$. If $\gamma \in \Gamma$, then $(\gamma')^{-1} = (\gamma^{-1})'$, and if $\gamma_1, \gamma_2 \in \Gamma$, then $\gamma_1'\gamma_2' = (\gamma_1\gamma_2)'$. Hence Γ' is a subgroup of $\mathrm{Isom}(V^\perp)$. The map $\mathrm{P}' : \Gamma \to \Gamma'$ defined by $\mathrm{P}'(\gamma) = \gamma'$ is epimorphism with kernel N, since N is a complete normal subgroup of Γ. Hence P' induces an isomorphism $\mathrm{P} : \Gamma/\mathrm{N} \to \Gamma'$ defined by $\mathrm{P}(\mathrm{N}\gamma) = \gamma'$. The group Γ' is a space group of isometries of $V^\perp$ with $V^\perp/\Gamma' = V^\perp/(\Gamma/\mathrm{N})$.

Let $\overline{\mathrm{N}} = \{\overline{\nu} : \nu \in \mathrm{N}\}$. Then $\overline{\mathrm{N}}$ is a subgroup of $\mathrm{Isom}(V)$. The map $\mathrm{B} : \mathrm{N} \to \overline{\mathrm{N}}$ defined by $\mathrm{B}(\nu) = \overline{\nu}$ is an isomorphism. The group $\overline{\mathrm{N}}$ is a space group of isometries of V with $V/\overline{\mathrm{N}} = V/\mathrm{N}$.

The action of Γ on V induces an action of Γ/N on V/N defined by

$$(\mathrm{N}\gamma)(\mathrm{N}v) = \mathrm{N}\overline{\gamma}v.$$

The action of Γ/N on V/N determines a homomorphism

$$\Xi : \Gamma/\mathrm{N} \to \mathrm{Isom}(V/\mathrm{N})$$

defined by $\Xi(\mathrm{N}\gamma) = \overline{\gamma}_\star$, where $\overline{\gamma}_\star : V/\mathrm{N} \to V/\mathrm{N}$ is defined by $\overline{\gamma}_\star(\mathrm{N}v) = \mathrm{N}\overline{\gamma}(v)$.

Theorem 2. *Let* M *be an* m*-space group, let* Δ *be an* $(n-m)$*-space group, and let* $\Theta : \Delta \to \mathrm{Isom}(E^m/\mathrm{M})$ *be a homomorphism. Identify* E^n *with* $E^m \times E^{n-m}$*, and extend* M *to a subgroup* N *of* $\mathrm{Isom}(E^n)$ *such that the point group of* N *acts trivially on* $(E^m)^\perp = E^{n-m}$*. Then there exists a unique* n*-space group* Γ *containing* N *as a complete normal subgroup such that* $\Gamma' = \Delta$*, and if* $\Xi : \Gamma/\mathrm{N} \to \mathrm{Isom}(E^m/\mathrm{N})$ *is the homomorphism induced by the action of* Γ/N *on* E^m/N*, then* $\Xi = \Theta\mathrm{P}$ *where* $\mathrm{P} : \Gamma/\mathrm{N} \to \Gamma'$ *is the isomorphism defined by* $\mathrm{P}(\mathrm{N}\gamma) = \gamma'$ *for each* $\gamma \in \Gamma$*.*

Proof. Let $\delta \in \Delta$. By Lemma 1 of [9], there exists $\tilde{\delta} \in N_E(\mathrm{M})$ such that $\tilde{\delta}_\star = \Theta(\delta)$. The isometry $\tilde{\delta}$ is unique up to multiplication by an element of M. Let $\delta = d + D$, with $d \in E^{n-m}$ and $D \in \mathrm{O}(n-m)$, and let $\tilde{\delta} = \tilde{d} + \tilde{D}$, with $\tilde{d} \in E^m$ and $\tilde{D} \in \mathrm{O}(m)$. Let $\hat{d} = \tilde{d} + d$, and let $\hat{D} = \tilde{D} \times D$. Then $\hat{D}E^m = E^m$ and $\hat{D}E^{n-m} = E^{n-m}$. Let $\hat{\delta} = \hat{d} + \hat{D}$. Then $\hat{\delta}$ is an isometry of E^n such that $\overline{\hat{\delta}} = \tilde{\delta}$ and $\big(\hat{\delta}\big)' = \delta$. The isometry $\hat{\delta}$ is unique up to multiplication by an element of N. We have that

$$\hat{\delta}^{-1} = -\hat{D}^{-1}\hat{d} + \hat{D}^{-1} = -\tilde{D}^{-1}\tilde{d} - D^{-1}d + \tilde{D}^{-1} \times D^{-1},$$

and so $\overline{\hat{\delta}^{-1}} = \big(\tilde{\delta}\big)^{-1}$ and $(\hat{\delta}^{-1})' = \delta^{-1}$. We have that

$$\overline{\hat{\delta}\mathrm{N}\hat{\delta}^{-1}} = \overline{\hat{\delta}}\,\overline{\mathrm{N}}\,\overline{\hat{\delta}^{-1}} = \tilde{\delta}\mathrm{M}\tilde{\delta}^{-1} = \mathrm{M} = \overline{\mathrm{N}}$$

and

$$(\hat{\delta}\mathrm{N}\hat{\delta}^{-1})' = \hat{\delta}'\mathrm{N}'(\hat{\delta}^{-1})' = \delta\{I'\}\delta^{-1} = \{I'\}.$$

Therefore $\hat{\delta}\mathrm{N}\hat{\delta}^{-1} = \mathrm{N}$.

Let Γ be the subgroup of $\mathrm{Isom}(E^n)$ generated by $\mathrm{N} \cup \{\hat{\delta} : \delta \in \Delta\}$. Then Γ contains N as a normal subgroup, and the point group of Γ leaves E^m invariant. Suppose $\gamma \in \Gamma$. Then there exists $\nu \in \mathrm{N}$ and $\delta_1, \dots, \delta_k \in \Delta$ and $\epsilon_1, \dots, \epsilon_k \in \{\pm 1\}$ such that $\gamma = \nu\hat{\delta}_1^{\epsilon_1} \cdots \hat{\delta}_k^{\epsilon_k}$. Then we have

$$\gamma' = \nu'(\hat{\delta}_1^{\epsilon_1})' \cdots (\hat{\delta}_k^{\epsilon_k})' = \delta_1^{\epsilon_1} \cdots \delta_k^{\epsilon_k}.$$

Hence $\Gamma' = \Delta$, and we have an epimorphism $\mathrm{P}' : \Gamma \to \Delta$ defined by $\mathrm{P}'(\gamma) = \gamma'$. The group N is in the kernel of P', and so P' induces an epimorphism $\mathrm{P} : \Gamma/\mathrm{N} \to \Delta$ defined by $\mathrm{P}(\mathrm{N}\gamma) = \gamma'$. Suppose $\mathrm{P}(\mathrm{N}\gamma) = I'$. By Lemma 1 of [9], we have that

$$\begin{aligned}
\mathrm{P}(\mathrm{N}\gamma) = I' \quad \Rightarrow \quad \gamma' = I' \quad &\Rightarrow \quad \delta_1^{\epsilon_1} \cdots \delta_k^{\epsilon_k} = I' \\
&\Rightarrow \quad \Theta(\delta_1)^{\epsilon_1} \cdots \Theta(\delta_k)^{\epsilon_k} = \overline{I}_\star \\
&\Rightarrow \quad (\tilde{\delta}_1)_\star^{\epsilon_1} \cdots (\tilde{\delta}_k)_\star^{\epsilon_k} = \overline{I}_\star \\
&\Rightarrow \quad \mathrm{M}(\tilde{\delta}_1)^{\epsilon_1} \cdots \mathrm{M}(\tilde{\delta}_k)^{\epsilon_k} = \mathrm{M} \\
&\Rightarrow \quad \mathrm{M}(\tilde{\delta}_1)^{\epsilon_1} \cdots (\tilde{\delta}_k)^{\epsilon_k} = \mathrm{M} \\
&\Rightarrow \quad \mathrm{M}\overline{\hat{\delta}_1^{\epsilon_1}} \cdots \overline{\hat{\delta}_k^{\epsilon_k}} = \mathrm{M} \quad \Rightarrow \quad \mathrm{M}\overline{\gamma} = \mathrm{M} \quad \Rightarrow \quad \overline{\gamma} \in \mathrm{M}.
\end{aligned}$$

As $\gamma' = I'$ and $\overline{\gamma} \in \overline{\mathrm{N}}$, we have that $\gamma \in \mathrm{N}$. Thus P is an isomorphism.

We next show that Γ acts discontinuously on E^n. Let C be a compact subset of E^n. Let K and L be the orthogonal projections of C into E^m and E^{n-m}, respectively. Then $C \subseteq K \times L$. As Δ acts discontinuously on E^{n-m}, there exists only finitely many elements $\delta_1, \dots, \delta_k$ of Δ such that $L \cap \delta_i L \neq \emptyset$ for each i. Let $\gamma_i = \hat{\delta}_i$ for $i = 1, \dots, k$. The set $K_i = K \cup \overline{\gamma}_i(K)$ is compact for each $i = 1, \dots, k$. As N acts discontinuously on E^m, there is a finite subset F_i of N such that $K_i \cap \nu K_i \neq \emptyset$ only if $\nu \in F_i$ for each i. Hence $K \cap \nu\overline{\gamma}_i(K) \neq \emptyset$ only if $\nu \in F_i$ for each i. The set $F = F_1\gamma_1 \cup \cdots \cup F_k\gamma_k$ is a finite subset of Γ. Suppose $\gamma \in \Gamma$ and $C \cap \gamma C \neq \emptyset$. Then $L \cap \gamma' L \neq \emptyset$, and so $\gamma' = \delta_i$ for some i. Hence $\gamma = \nu\gamma_i$ for some $\nu \in \mathrm{N}$. Now we have that $K \cap \overline{\gamma}K \neq \emptyset$, and so $K \cap \nu\overline{\gamma}_i K \neq \emptyset$. Hence $\nu \in F_i$. Therefore $\gamma \in F$. Thus Γ acts discontinuously on E^n. Therefore Γ is a discrete subgroup of $\mathrm{Isom}(E^n)$ by Theorem 5.3.5 of [7].

Let D_{M} and D_Δ be fundamental domains for M in E^m and Δ in E^{n-m}, respectively. Then their topological closures $\overline{D}_{\mathrm{M}}$ and $\overline{D}_\Delta$ are compact sets. Let $x \in E^n$. Write $x = \overline{x} + x'$ with $\overline{x} \in E^m$ and $x' \in E^{n-m}$. Then there exists $\delta \in \Delta$ such that $\delta x' \in \overline{D}_\Delta$, and there exists $\nu \in \mathrm{N}$ such that $\nu\tilde{\delta}\overline{x} \in \overline{D}_{\mathrm{M}}$. We have that

$$\nu\hat{\delta}x = \overline{\nu\hat{\delta}\overline{x}} + (\nu\hat{\delta})'x' = \nu\tilde{\delta}\overline{x} + \delta x' \in \overline{D}_{\mathrm{M}} \times \overline{D}_\Delta.$$

Hence the quotient map $\pi : E^n \to E^n/\Gamma$ maps the compact set $\overline{D}_{\mathrm{M}} \times \overline{D}_\Delta$ onto E^n/Γ. Therefore E^n/Γ is compact. Thus Γ is an n-space group.

Let $\Xi : \Gamma/\mathrm{N} \to \mathrm{Isom}(E^n/\mathrm{N})$ be the homomorphism induced by the action of Γ/N on E^m/N. Let $\gamma \in \Gamma$. Then there exists $\delta \in \Delta$ such that $\mathrm{N}\gamma = \mathrm{N}\hat{\delta}$. Then $\gamma' = (\hat{\delta})' = \delta$, and we have that $\Xi = \Theta\mathrm{P}$, since

$$\Xi(\mathrm{N}\gamma) = \Xi(\mathrm{N}\hat{\delta}) = (\overline{\hat{\delta}})_\star = \tilde{\delta}_\star = \Theta(\delta) = \Theta(\gamma') = \Theta\mathrm{P}(\mathrm{N}\gamma).$$

Suppose γ is an isometry of E^n such that $\gamma\mathrm{N}\gamma^{-1} = \mathrm{N}$ and $\gamma' \in \Delta$ and $\overline{\gamma}_\star = \Theta(\gamma')$. Then $\widehat{\gamma'} \in \Gamma$. Now $\overline{\gamma}_\star = \widetilde{\gamma'}_\star$, and so $\overline{\gamma} = \overline{\nu}\widetilde{\gamma'}$ for some $\nu \in \mathrm{N}$ by Lemma 1 of [9]. Then $\gamma = \nu\widehat{\gamma'}$. Hence $\gamma \in \Gamma$. Thus Γ is the unique n-space group that contains N as a complete normal subgroup such that $\Gamma' = \Delta$ and $\Xi = \Theta\mathrm{P}$. □

3. Isomorphisms of Pairs of Space Groups

An *affinity* α of E^n is a map $\alpha : E^n \to E^n$ for which there is an element $a \in E^n$ and a matrix $A \in \mathrm{GL}(n, \mathbb{R})$ such that $\alpha(x) = a + Ax$ for all x in E^n. We write simply $\alpha = a + A$. The set $\mathrm{Aff}(E^n)$ of all affinities of E^n is a group that contains $\mathrm{Isom}(E^n)$ as a subgroup.

Let N_i be a complete normal subgroup of an n-space group Γ_i for $i = 1, 2$. We want to know when (Γ_1, N_1) is isomorphic to (Γ_2, N_2), that is, when there is an isomorphism $\zeta : \Gamma_1 \to \Gamma_2$ such that $\zeta(\mathrm{N}_1) = \mathrm{N}_2$. By a Theorem of Bieberbach an isomorphism $\zeta : \Gamma_1 \to \Gamma_2$ is equal to conjugation by an affinity ϕ of E^n. In this section, we determine necessary and sufficient conditions such that there exists an affinity ϕ of E^n such that $\phi(\Gamma_1, \mathrm{N}_1)\phi^{-1} = (\Gamma_2, \mathrm{N}_2)$.

Let ϕ be an affinity of E^n such that $\phi(\Gamma_1, \mathrm{N}_1)\phi^{-1} = (\Gamma_2, \mathrm{N}_2)$. Write $\phi = c + C$ with $c \in E^n$ and $C \in \mathrm{GL}(n, \mathbb{R})$. Let $V_i = \mathrm{Span}(\mathrm{N}_i)$, for $i = 1, 2$. Let $a + I \in \mathrm{N}_1$. Then $\phi(a + I)\phi^{-1} = Ca + I$. Hence $CV_1 \subseteq V_2$. Let $\overline{C} : V_1 \to V_2$ be the linear transformation obtained by restricting C. Let $\overline{C'} : V_1^{\perp} \to V_2$ and $C' : V_1^{\perp} \to V_2^{\perp}$ be the linear transformations obtained by restricting C to $V_1^{\perp}$ followed by the orthogonal projections to V_2 and $V_2^{\perp}$, respectively. Write $c = \overline{c} + c'$ with $\overline{c} \in V_2$ and $c' \in V_2^{\perp}$. Let $\overline{\phi} : V_1 \to V_2$ and $\phi' : V_1^{\perp} \to V_2^{\perp}$ be the affine transformations defined by $\overline{\phi} = \overline{c} + \overline{C}$ and $\phi' = c' + C'$.

Lemma 2. *Let N_i be a complete normal subgroup of an n-space group Γ_i, with $V_i = \mathrm{Span}(\mathrm{N}_i)$, for $i = 1, 2$. Let $\phi = c + C$ be an affinity of E^n such that $\phi(\Gamma_1, \mathrm{N}_1)\phi^{-1} = (\Gamma_2, \mathrm{N}_2)$. Then $\overline{C}$ and C' are invertible, with $\overline{C}^{-1} = \overline{C^{-1}}$ and $(C')^{-1} = (C^{-1})'$, and $\overline{(C^{-1})'} = -\overline{C}^{-1}\overline{C'}(C')^{-1}$. If $b + B \in \Gamma_1$, then $\overline{C'}B' = \overline{CBC}^{-1}\overline{C'}$. Moreover $\overline{C'}V_1^{\perp} \subseteq \mathrm{Span}(Z(\mathrm{N}_2))$.*

Proof. We have that

$$\dim V_1 = \dim \mathrm{N}_1 = \dim \mathrm{N}_2 = \dim V_2.$$

Let $y \in E^n$ and write $y = \overline{y} + y'$ with $\overline{y} \in V_2$ and $y' \in V_2^{\perp}$. Then

$$\overline{y} = CC^{-1}(\overline{y}) = \overline{C}\,\overline{C^{-1}}(\overline{y}),$$

and so $\overline{C}$ is invertible with $\overline{C}^{-1} = \overline{C^{-1}}$. We have that

$$\begin{aligned} y' &= CC^{-1}y' \\ &= C(\overline{(C^{-1})'}(y') + (C^{-1})'(y')) \\ &= C\overline{(C^{-1})'}(y') + C(C^{-1})'(y') \\ &= \overline{C}\,\overline{(C^{-1})'}(y') + \overline{C'}(C^{-1})'(y') + C'(C^{-1})'(y'). \end{aligned}$$

Hence C' is invertible, with $(C')^{-1} = (C^{-1})'$, and $\overline{C}\,\overline{(C^{-1})'} + \overline{C'}(C^{-1})' = 0$. Therefore $\overline{(C^{-1})'} = -\overline{C}^{-1}\overline{C'}(C')^{-1}$.

Let $\gamma = b + B \in \Gamma_1$. Then $\phi\gamma\phi^{-1} \in \Gamma_2$. Let $w \in V_2^\perp$. Then we have

$$\begin{aligned} w &= CBC^{-1}w \\ &= CB(\overline{(C^{-1})'}(w) + (C^{-1})'(w)) \\ &= C(\overline{B(C^{-1})'}(w) + B'(C^{-1})'(w)) \\ &= C\overline{B(C^{-1})'}(w) + CB'(C^{-1})'(w) \\ &= \overline{CB(C^{-1})'}(w) + \overline{C'}B'(C^{-1})'(w) + C'B'(C^{-1})'(w). \end{aligned}$$

As $\phi\gamma\phi^{-1}(w) \in V_2^\perp$, we have that

$$\overline{CB(C^{-1})'} + \overline{C'}B'(C^{-1})' = 0.$$

Therefore

$$\overline{CB}(-\overline{C}^{-1}\overline{C'}(C')^{-1}) + \overline{C'}B'(C')^{-1} = 0.$$

Hence $\overline{C'}B' = \overline{CBC}^{-1}\overline{C'}$.

Now suppose $\gamma \in \mathrm{N}_1$. Then $B' = I'$ by Theorem 1(2). Hence $\overline{BC}^{-1}\overline{C'} = \overline{C}^{-1}\overline{C'}$. By Lemma 5 of [9], we deduce that $\overline{C}^{-1}\overline{C'}V_1^\perp \subseteq \mathrm{Span}(Z(\mathrm{N}_1))$. As $\phi Z(\mathrm{N}_1)\phi^{-1} = Z(\mathrm{N}_2)$, we have that $\overline{C'}V_1^\perp \subseteq \mathrm{Span}(Z(\mathrm{N}_2))$. □

Theorem 3. *Let N_i be a complete normal subgroup of an n-space group Γ_i, with $V_i = \mathrm{Span}(\mathrm{N}_i)$ for $i = 1, 2$. Let $\Xi_i : \Gamma_i/\mathrm{N}_i \to \mathrm{Isom}(V_i/\mathrm{N}_i)$ be the homomorphism induced by the action of Γ_i/N_i on V_i/N_i for $i = 1, 2$. Let $\alpha : V_1 \to V_2$, and $\beta : V_1^\perp \to V_2^\perp$ be affinities such that $\alpha\overline{\mathrm{N}}_1\alpha^{-1} = \overline{\mathrm{N}}_2$ and $\beta\Gamma_1'\beta^{-1} = \Gamma_2'$. Let $\alpha_\star : V_1/\mathrm{N}_1 \to V_2/\mathrm{N}_2$ be the affinity defined by $\alpha_\star(\mathrm{N}_1 v) = \mathrm{N}_2\alpha(v)$, and let $\alpha_\sharp : \mathrm{Aff}(V_1/\mathrm{N}_1) \to \mathrm{Aff}(V_2/\mathrm{N}_2)$ be the isomorphism defined by $\alpha_\sharp(\chi) = \alpha_\star\chi\alpha_\star^{-1}$. Let $\beta_* : \Gamma_1' \to \Gamma_2'$ be the isomorphism defined by $\beta_*(\gamma') = \beta\gamma'\beta^{-1}$.*

Write $\alpha = \overline{c} + \overline{C}$ with $\overline{c} \in V_2$ and $\overline{C} : V_1 \to V_2$ a linear isomorphism. Let $D : V_1^\perp \to \mathrm{Span}(Z(\mathrm{N}_2))$ be a linear transformation such that if $b + B \in \Gamma_1$, then $DB' = \overline{CBC}^{-1}D$. Let $\mathrm{P}_i : \Gamma_i/\mathrm{N}_i \to \Gamma_i'$ be the isomorphism defined by $\mathrm{P}_i(\mathrm{N}_i\gamma) = \gamma'$ for each $i = 1, 2$, and let $p_1 : \Gamma_1/\mathrm{N}_1 \to V_1^\perp$ be defined by $p_1(\mathrm{N}_1(b+B)) = b'$. Let $\mathcal{K}_i$ be the connected component of the identity of the Lie group $\mathrm{Isom}(V_i/\mathrm{N}_i)$ for each $i = 1, 2$. Note that $\mathcal{K}_i = \{(v + \overline{I})_\star : v \in \mathrm{Span}(Z(\mathrm{N}_i))\}$ by Theorem 1 of [9]. *Then the following are equivalent:*

(1) *There exists an affinity $\phi = c + C$ of E^n such that $\phi(\Gamma_1, \mathrm{N}_1)\phi^{-1} = (\Gamma_2, \mathrm{N}_2)$, with $\overline{\phi} = \alpha$, and $\phi' = \beta$, and $\overline{C'} = D$.*

(2) *We have that*

$$\Xi_2\mathrm{P}_2^{-1}\beta_*\mathrm{P}_1 = (Dp_1)_\star\alpha_\sharp\Xi_1$$

with $(Dp_1)_\star : \Gamma_1/\mathrm{N}_1 \to \mathcal{K}_2$ a crossed homomorphism defined by

$$(Dp_1)_\star(\mathrm{N}_1(b + B)) = (Db' + \overline{I})_\star$$

and Γ_1/N_1 acting on $\mathcal{K}_2$ by $\mathrm{N}_1(b + B)(v + \overline{I})_\star = (\overline{CBC}^{-1}v + \overline{I})_\star$ for each $b + B \in \Gamma_1$ and $v \in \mathrm{Span}(Z(\mathrm{N}_2))$.

(3) *We have that*

$$\alpha_\sharp^{-1}\Xi_2\mathrm{P}_2^{-1}\beta_*\mathrm{P}_1 = (\overline{C}^{-1}Dp_1)_\star\Xi_1$$

with $(\overline{C}^{-1}Dp_1)_\star : \Gamma_1/\mathrm{N}_1 \to \mathcal{K}_1$ a crossed homomorphism defined by

$$(\overline{C}^{-1}Dp_1)_\star(\mathrm{N}_1(b + B)) = (\overline{C}^{-1}D(b') + \overline{I})_\star$$

and Γ_1/N_1 *acting on* $\mathcal{K}_1$ *by* $\mathrm{N}_1(b+B)(v+\overline{I})_\star = (\overline{B}v+\overline{I})_\star$ *for each* $b+B \in \Gamma_1$ *and* $v \in \mathrm{Span}(Z(\mathrm{N}_1))$.

Proof. Suppose there exists an affinity $\phi = c + C$ of E^n such that $\phi(\Gamma_1, \mathrm{N}_1)\phi^{-1} = (\Gamma_2, \mathrm{N}_2)$, with $\overline{\phi} = \alpha$, and $\phi' = \beta$, and $\overline{C'} = D$. Let $\gamma = b + B \in \Gamma_1$. Then we have

$$\phi\gamma\phi^{-1} = (c+C)(b+B)(c+C)^{-1} = Cb + (I - CBC^{-1})c + CBC^{-1}.$$

Hence we have

$$\begin{aligned}\overline{\phi\gamma\phi^{-1}} &= \overline{Cb} + \overline{C'}b' + (\overline{I} - \overline{CBC}^{-1})\overline{c} + \overline{CBC}^{-1} \\ &= (\overline{C'}b' + \overline{I})(\overline{c} + \overline{C})(\overline{b} + \overline{B})(\overline{c} + \overline{C})^{-1} \\ &= (Db' + \overline{I})\overline{\phi}\overline{\gamma}\overline{\phi}^{-1} \quad = \quad (Db' + \overline{I})\alpha\overline{\gamma}\alpha^{-1},\end{aligned}$$

and

$$\begin{aligned}(\phi\gamma\phi^{-1})' &= C'b' + (I' - C'B'(C')^{-1})c' + C'B'(C')^{-1} \\ &= (c' + C')(b' + B')(c' + C')^{-1} \\ &= \phi'\gamma'(\phi')^{-1} \quad = \quad \beta\gamma'\beta^{-1}.\end{aligned}$$

Observe that

$$\begin{aligned}\Xi_2\mathrm{P}_2^{-1}\beta_*\mathrm{P}_1(\mathrm{N}_1\gamma) &= \Xi_2\mathrm{P}_2^{-1}\beta_*(\gamma') \\ &= \Xi_2\mathrm{P}_2^{-1}(\beta\gamma'\beta^{-1}) \\ &= \Xi_2\mathrm{P}_2^{-1}(\phi\gamma\phi^{-1})' \\ &= \Xi_2(\mathrm{N}_2\phi\gamma\phi^{-1}) \\ &= (\overline{\phi\gamma\phi^{-1}})_\star \\ &= ((Db' + \overline{I})\alpha\overline{\gamma}\alpha^{-1})_\star \\ &= (Db' + \overline{I})_\star\alpha_\star\overline{\gamma}_\star\alpha_\star^{-1} \\ &= (Dp_1(\mathrm{N}_1(b+B)))_\star\alpha_\sharp(\overline{\gamma}_\star) \quad = \quad (Dp_1(\mathrm{N}_1\gamma))_\star\alpha_\sharp(\Xi_1(\mathrm{N}_1\gamma)).\end{aligned}$$

Hence we have that $\Xi_2\mathrm{P}_2^{-1}\beta_*\mathrm{P}_1 = (Dp_1)_\star\alpha_\sharp\Xi_1$.

Conversely, suppose that $\Xi_2\mathrm{P}_2^{-1}\beta_*\mathrm{P}_1 = (Dp_1)_\star\alpha_\sharp\Xi_1$. Define an affine transformation $\phi : E^n \to E^n$ by

$$\phi(x) = \alpha(\overline{x}) + D(x') + \beta(x')$$

for each $x \in E^n$, where $x = \overline{x} + x'$ with $\overline{x} \in V_1$ and $x' \in V_1^\perp$. Then ϕ is an affinity of E^n, with

$$\phi^{-1}(y) = \alpha^{-1}(\overline{y}) - \overline{C}^{-1}D\beta^{-1}(y') + \beta^{-1}(y')$$

for each $y \in E^n$ where $y = \overline{y} + y'$ with $\overline{y} \in V_2$ and $y' \in V_2^\perp$.

Write $\beta = c' + C'$ with $c' \in V_2^\perp$ and $C' : V_1^\perp \to V_2^\perp$ a linear isomorphism. Write $\phi = c + C$ with $c \in E^n$ and C a linear isomorphism of E^n. Then $c = \overline{c} + c'$ and $Cx = \overline{C}\overline{x} + Dx' + C'x'$ for each $x \in E^n$ with $\overline{x} \in V_1$ and $x' \in V_1^\perp$. We have that $\overline{\phi} = \alpha$, and $\phi' = \beta$ and $\overline{C'} = D$. We also have $\phi^{-1} = -C^{-1}c + C^{-1}$ and

$$C^{-1}y = \overline{C}^{-1}\overline{y} - \overline{C}^{-1}D(C')^{-1}y' + (C')^{-1}y'$$

for all $y \in E^n$ with $\overline{y} \in V_2$ and $y' \in V_2^\perp$.

Let $\gamma \in \Gamma_1$. Write $\gamma = b + B$ with $b \in E^n$ and $B \in \mathrm{O}(n)$. Then we have

$$\phi\gamma\phi^{-1} = (c+C)(b+B)(c+C)^{-1} = Cb + (I - CBC^{-1})c + CBC^{-1}.$$

Let $y \in E^n$. Write $y = \overline{y} + y'$ with $\overline{y} \in V_2$ and $y' \in V_2^\perp$. Then we have that

$$\begin{aligned} CBC^{-1} &= CB(\overline{C}^{-1}\overline{y} - \overline{C}^{-1}D(C')^{-1}y' + (C')^{-1}y' \\ &= C(\overline{BC}^{-1}\overline{y} - \overline{BC}^{-1}D(C')^{-1}y' + B'(C')^{-1}y' \\ &= \overline{CBC}^{-1}\overline{y} - \overline{CBC}^{-1}D(C')^{-1}y' + DB'(C')^{-1}y' + C'B'(C')^{-1}y' \\ &= \overline{CBC}^{-1}\overline{y} + C'B'(C')^{-1}y'. \end{aligned}$$

Hence $CBC^{-1} = \overline{CBC}^{-1} \times C'B'(C')^{-1}$ as a linear isomorphism of $E^n = V_2 \times V_2^\perp$.

Moreover, we have

$$\begin{aligned} \overline{\phi\gamma\phi^{-1}} &= \overline{C}\overline{b} + \overline{C}'b' + (\overline{I} - \overline{CBC}^{-1})\overline{c} + \overline{CBC}^{-1} \\ &= (\overline{C}'b' + \overline{I})(\overline{c} + \overline{C})(\overline{b} + \overline{B})(\overline{c} + \overline{C})^{-1} \\ &= (Db' + \overline{I})\overline{\phi}\overline{\gamma}\overline{\phi}^{-1} \quad = \quad (Db' + \overline{I})\alpha\overline{\gamma}\alpha^{-1}, \end{aligned}$$

and

$$\begin{aligned} (\phi\gamma\phi^{-1})' &= C'b' + (I' - C'B'(C')^{-1})c' + C'B'(C')^{-1} \\ &= (c' + C')(b' + B')(c' + C')^{-1} \\ &= \phi'\gamma'(\phi')^{-1} \quad = \quad \beta\gamma'\beta^{-1}. \end{aligned}$$

As $\Xi_2 \mathrm{P}_2^{-1}\beta_*\mathrm{P}_1 = (Dp_1)_\star\alpha_\sharp\Xi_1$, we have that $(\alpha\overline{\gamma}\alpha^{-1})_\star$ is an isometry of V_2/N_2. By Lemmas 1 and 7 of [9], we have that $\alpha\overline{\gamma}\alpha^{-1}$ is an isometry of V_2. Hence $\overline{CBC}^{-1}$ is an orthogonal transformation of V_2.

As $\beta\Gamma_1'\beta^{-1} = \Gamma_2'$, we have that $C'B'(C')^{-1}$ is an orthogonal transformation of $V_2^\perp$. Hence $CBC^{-1} = \overline{CBC}^{-1} \times C'B'(C')^{-1}$ is an orthogonal transformation of $E^n = V_2 \times V_2^\perp$. Therefore $\phi\gamma\phi^{-1}$ is an isometry of E^n for each $\gamma \in \Gamma_1$.

As Γ_1 acts discontinuously on E^n and ϕ is a homeomorphism of E^n, we have that $\phi\Gamma_1\phi^{-1}$ acts discontinuously on E^n. Therefore $\phi\Gamma_1\phi^{-1}$ is a discrete subgroup of $\mathrm{Isom}(E^n)$ by Theorem 5.3.5 of [7]. Now ϕ induces a homeomorphism $\phi_\star : E^n/\Gamma_1 \to E^n/\phi\Gamma_1\phi^{-1}$ defined by $\phi_\star(\Gamma_1 x) = \phi\Gamma_1\phi^{-1}\phi(x)$. Hence $E^n/\phi\Gamma_1\phi^{-1}$ is compact. Therefore $\phi\Gamma_1\phi^{-1}$ is a n-space group.

Now $\phi_* : \Gamma_1 \to \phi\Gamma_1\phi^{-1}$, defined by $\phi_*(\gamma) = \phi\gamma\phi^{-1}$, is an isomorphism that maps the normal subgroup N_1 to the normal subgroup $\phi\mathrm{N}_1\phi^{-1}$ of $\phi\Gamma_1\phi^{-1}$, and $\phi\Gamma_1\phi^{-1}/\phi\mathrm{N}_1\phi^{-1}$ is isomorphic to Γ_1/N_1. Hence $\phi\Gamma_1\phi^{-1}/\phi\mathrm{N}_1\phi^{-1}$ is a space group. Therefore $\phi\mathrm{N}_1\phi^{-1}$ is a complete normal subgroup of $\phi\Gamma_1\phi^{-1}$ by Theorem 5 of [8].

Now suppose $\nu = a + A \in \mathrm{N}_1$. Then $a' = 0$ and $A' = I'$, and so $\nu' = I'$. Hence $\overline{\phi\nu\phi^{-1}} = \alpha\overline{\nu}\alpha^{-1}$ and $(\phi\nu\phi^{-1})' = I'$. As $\alpha\overline{\mathrm{N}}_1\alpha^{-1} = \overline{\mathrm{N}}_2$, we have that $\phi\mathrm{N}_1\phi^{-1} = \mathrm{N}_2$. Moreover, as $\beta\Gamma_1'\beta^{-1} = \Gamma_2'$, we have that $(\phi\Gamma_1\phi^{-1})' = \Gamma_2'$.

Let $\Xi : \phi\Gamma_1\phi^{-1}/\mathrm{N}_2 \to \mathrm{Isom}(V_2/\mathrm{N}_2)$ be the homomorphism induced by the action of $\phi\Gamma_1\phi^{-1}/\mathrm{N}_2$ on V_2/N_2. Let $\gamma = b + B \in \Gamma_1$, and let $\mathrm{P} : \phi\Gamma_1\phi^{-1}/\mathrm{N}_2 \to \Gamma_2'$ be the

isomorphism defined by $\mathrm{P}(\mathrm{N}_2\phi\gamma\phi^{-1}) = (\phi\gamma\phi^{-1})'$. Then we have that

$$\begin{aligned}
\Xi\mathrm{P}^{-1}(\beta\gamma'\beta^{-1}) &= \Xi\mathrm{P}^{-1}((\phi\gamma\phi^{-1})') \\
&= \Xi(\mathrm{N}_2\phi\gamma\phi^{-1}) \\
&= (\overline{\phi\gamma\phi^{-1}})_\star \\
&= ((Db' + \overline{I})\alpha\overline{\gamma}\alpha^{-1})_\star \\
&= (Db' + \overline{I})_\star\alpha_\star\overline{\gamma}_\star\alpha_\star^{-1} \\
&= (Dp_1(\mathrm{N}_1(b+B)))_\star\alpha_\sharp(\overline{\gamma}_\star) \\
&= (Dp_1(\mathrm{N}_1\gamma))_\star\alpha_\sharp(\Xi_1(\mathrm{N}_1\gamma)) \\
&= \Xi_2\mathrm{P}_2^{-1}\beta_*\mathrm{P}_1(\mathrm{N}_1\gamma) \quad = \quad \Xi_2\mathrm{P}_2^{-1}(\beta\gamma'\beta^{-1}).
\end{aligned}$$

Hence we have that $\Xi\mathrm{P}^{-1} = \Xi_2\mathrm{P}_2^{-1}$. Therefore $\phi\Gamma_1\phi^{-1} = \Gamma_2$ by Theorem 2. Thus $\phi(\Gamma_1, \mathrm{N}_1)\phi^{-1} = (\Gamma_2, \mathrm{N}_2)$.

Let $\gamma = b + B$ and $\gamma_1 = b_1 + B_1$ be elements of Γ_1. Then we have that

$$\begin{aligned}
(Dp_1)_\star(\mathrm{N}_1\gamma\mathrm{N}_1\gamma_1) &= (Dp_1)_\star(\mathrm{N}_1(b + Bb_1 + BB_1)) \\
&= (D(b + Bb_1)' + \overline{I})_\star \\
&= (D(b' + B'b_1') + \overline{I})_\star \\
&= (Db' + DB'b_1' + \overline{I})_\star \\
&= (Db' + \overline{CBC}^{-1}Db_1' + \overline{I})_\star \\
&= (Db' + \overline{I})_\star(\overline{CBC}^{-1}Db_1' + \overline{I})_\star \\
&= (Db' + \overline{I})_\star(\mathrm{N}_1(b+B))(Db_1' + \overline{I})_\star \\
&= (Dp_1)_\star(\mathrm{N}_1\gamma)((\mathrm{N}_1\gamma)(Dp_1)_\star(\mathrm{N}_1\gamma_1)).
\end{aligned}$$

Therefore $(Dp_1)_\star : \Gamma_1/\mathrm{N}_1 \to \mathcal{K}_2$ is a crossed homomorphism. Thus statements (1) and (2) are equivalent.

The equation $\Xi_2\mathrm{P}_2^{-1}\beta_*\mathrm{P}_1 = (Dp_1)_\star\alpha_\sharp\Xi_1$ is equivalent to the equation

$$\alpha_\sharp^{-1}\Xi_2\mathrm{P}_2^{-1}\beta_*\mathrm{P}_1 = \alpha_\sharp^{-1}(Dp_1)_\star\alpha_\sharp\Xi_1.$$

Observe that

$$\begin{aligned}
\alpha_\sharp^{-1}(Dp_1)_\star\alpha_\sharp\Xi_1(\mathrm{N}_1\gamma) &= \alpha_\sharp^{-1}((Dp_1)_\star(\mathrm{N}_1\gamma)\alpha_\sharp\Xi_1(\mathrm{N}_1\gamma)) \\
&= \alpha_\sharp^{-1}((Db' + \overline{I})_\star\alpha_\sharp\overline{\gamma}_\star) \\
&= \alpha_\sharp^{-1}((Db' + \overline{I})_\star\alpha_\star\overline{\gamma}_\star\alpha_\star^{-1}) \\
&= \alpha_\star^{-1}(Db' + \overline{I})_\star\alpha_\star\overline{\gamma}_\star\alpha_\star^{-1}\alpha_\star \\
&= (\alpha^{-1}(Db' + \overline{I})\alpha)_\star\overline{\gamma}_\star \\
&= (\overline{C}^{-1}Db' + \overline{I})_\star\Xi_1(\mathrm{N}_1\gamma) \\
&= (\overline{C}^{-1}Dp_1)_\star(\mathrm{N}_1\gamma)\Xi_1(\mathrm{N}_1\gamma).
\end{aligned}$$

Hence we have that $\alpha_\sharp^{-1}(Dp_1)_\star\alpha_\sharp\Xi_1 = (\overline{C}^{-1}Dp_1)_\star\Xi_1$. By the same argument as with $(Dp_1)_\star : \Gamma_1/\mathrm{N}_1 \to \mathcal{K}_2$, we have that $(\overline{C}^{-1}Dp_1)_\star : \Gamma_1/\mathrm{N}_1 \to \mathcal{K}_1$ is a crossed homomorphism. Thus (2) and (3) are equivalent. □

4. Outer Automorphism Groups of Space Groups

Through this section, let m be a positive integer less than n. Let M be an m-space group and let Δ be an $(n-m)$-space group.

Definition: Define $\mathrm{Iso}(\Delta, \mathrm{M})$ to be the set of isomorphism classes of pairs (Γ, N) where N is a complete normal subgroup of an n-space group Γ such that N is isomorphic to M and Γ/N is isomorphic to Δ. We denote the isomorphism class of a pair (Γ, N) by $[\Gamma, \mathrm{N}]$.

Let N be a complete normal subgroup of an n-space group Γ, and let $\mathrm{Out}_E(\mathrm{N})$ be the Euclidean outer automorphism group of N defined in §4 of [9]. The group $\mathrm{Out}_E(\mathrm{N})$ is finite by Theorem 2 of [9]. The action of Γ on N by conjugation induces a homomorphism

$$\mathcal{O} : \Gamma/\mathrm{N} \to \mathrm{Out}_E(\mathrm{N})$$

defined by $\mathcal{O}(\mathrm{N}\gamma) = \gamma_*\mathrm{Inn}(\mathrm{N})$ where $\gamma_*(\nu) = \gamma\nu\gamma^{-1}$ for each $\gamma \in \Gamma$ and $\nu \in \mathrm{N}$. Let $\alpha : \mathrm{N}_1 \to \mathrm{N}_2$ be an isomorphism. Then α induces an isomorphism

$$\alpha_\# : \mathrm{Out}(\mathrm{N}_1) \to \mathrm{Out}(\mathrm{N}_2)$$

defined by $\alpha_\#(\zeta\mathrm{Inn}(\mathrm{N}_1)) = \alpha\zeta\alpha^{-1}\mathrm{Inn}(\mathrm{N}_2)$ for each $\zeta \in \mathrm{Aut}(\mathrm{N}_1)$.

Lemma 3. *Let N_i be a complete normal subgroup of an n-space group Γ_i for $i = 1, 2$. Let $\mathcal{O}_i : \Gamma_i/\mathrm{N}_i \to \mathrm{Out}_E(\mathrm{N}_i)$ be the homomorphism induced by the action of Γ_i on N_i by conjugation for $i = 1, 2$, and let $\alpha : \mathrm{N}_1 \to \mathrm{N}_2$ and $\phi : \Gamma_1 \to \Gamma_2$ and $\beta : \Gamma_1/\mathrm{N}_1 \to \Gamma_2/\mathrm{N}_2$ be isomorphisms such that the following diagram commutes*

$$\begin{array}{ccccccccc} 1 & \to & \mathrm{N}_1 & \to & \Gamma_1 & \to & \Gamma_1/\mathrm{N}_1 & \to & 1 \\ & & \downarrow \alpha & & \downarrow \phi & & \downarrow \beta & & \\ 1 & \to & \mathrm{N}_2 & \to & \Gamma_2 & \to & \Gamma_2/\mathrm{N}_2 & \to & 1, \end{array}$$

where the horizontal maps are inclusions and projections, then $\mathcal{O}_2 = \alpha_\#\mathcal{O}_1\beta^{-1}$.

Proof. Let $\gamma \in \Gamma_1$. Then we have that

$$\mathcal{O}_2(\mathrm{N}_2\phi(\gamma)) = \phi(\gamma)_*\mathrm{Inn}(\mathrm{N}_2),$$

whereas

$$\begin{aligned} \alpha_\#\mathcal{O}_1\beta^{-1}(\mathrm{N}_2\phi(\gamma)) &= \alpha_\#\mathcal{O}_1(\mathrm{N}_1\gamma) \\ &= \alpha_\#(\gamma_*\mathrm{Inn}(\mathrm{N}_1)) = \alpha\gamma_*\alpha^{-1}\mathrm{Inn}(\mathrm{N}_2). \end{aligned}$$

If $\nu \in \mathrm{N}$, then

$$\begin{aligned} \alpha\gamma_*\alpha^{-1}(\nu) &= \alpha\gamma_*\alpha^{-1}(\nu) \\ &= \alpha(\gamma\alpha^{-1}(\nu)\gamma^{-1}) \\ &= \phi(\gamma\phi^{-1}(\nu)\gamma^{-1}) \\ &= \phi(\gamma)\nu\phi(\gamma)^{-1} = \phi(\gamma)_*(\nu). \end{aligned}$$

Hence $\alpha\gamma_*\alpha^{-1} = \phi(\gamma)_*$. Therefore $\mathcal{O}_2 = \alpha_\#\mathcal{O}_1\beta^{-1}$. □

Definition: Define $\mathrm{Hom}_f(\Delta, \mathrm{Out}(\mathrm{M}))$ to be the set of all homomorphisms from Δ to $\mathrm{Out}(\mathrm{M})$ that have finite image.

The group $\mathrm{Out}(\mathrm{M})$ acts on the left of $\mathrm{Hom}_f(\Delta, \mathrm{Out}(\mathrm{M}))$ by conjugation, that is, if $g \in \mathrm{Out}(\mathrm{M})$ and $\eta \in \mathrm{Hom}_f(\Delta, \mathrm{Out}(\mathrm{M}))$, then $g\eta = g_*\eta$ where $g_* : \mathrm{Out}(\mathrm{M}) \to$

$\mathrm{Out}(\mathrm{M})$ is defined by $g_*(h) = ghg^{-1}$. Let $\mathrm{Out}(\mathrm{M})\backslash\mathrm{Hom}_f(\Delta, \mathrm{Out}(\mathrm{M}))$ be the set of $\mathrm{Out}(\mathrm{M})$-orbits. The group $\mathrm{Aut}(\Delta)$ acts on the right of $\mathrm{Hom}_f(\Delta, \mathrm{Out}(\mathrm{M}))$ by composition of homomorphisms. If $\beta \in \mathrm{Aut}(\Delta)$ and $\eta \in \mathrm{Hom}_f(\Delta, \mathrm{Out}(\mathrm{M}))$ and $g \in \mathrm{Out}(\mathrm{M})$, then

$$(g\eta)\beta = (g_*\eta)\beta = g_*(\eta\beta) = g(\eta\beta).$$

Hence $\mathrm{Aut}(\Delta)$ acts on the right of $\mathrm{Out}(\mathrm{M})\backslash\mathrm{Hom}_f(\Delta, \mathrm{Out}(\mathrm{M}))$ by

$$(\mathrm{Out}(\mathrm{M})\eta)\beta = \mathrm{Out}(\mathrm{M})(\eta\beta).$$

Let $\delta, \epsilon \in \Delta$ and $\eta \in \mathrm{Hom}_f(\Delta, \mathrm{Out}(\mathrm{M}))$. Then we have that

$$\eta\delta_*(\epsilon) = \eta(\delta\epsilon\delta^{-1}) = \eta(\delta)\eta(\epsilon)\eta(\delta)^{-1} = \eta(\delta)_*\eta(\epsilon) = (\eta(\delta)\eta)(\epsilon).$$

Hence $\eta\delta_* = \eta(\delta)\eta$. Therefore $\mathrm{Inn}(\Delta)$ acts trivially on $\mathrm{Out}(\mathrm{M})\backslash\mathrm{Hom}_f(\Delta, \mathrm{Out}(\mathrm{M}))$. Hence $\mathrm{Out}(\Delta)$ acts on the right of $\mathrm{Out}(\mathrm{M})\backslash\mathrm{Hom}_f(\Delta, \mathrm{Out}(\mathrm{M}))$ by

$$(\mathrm{Out}(\mathrm{M})\eta)(\beta\mathrm{Inn}(\Delta)) = \mathrm{Out}(\mathrm{M})(\eta\beta).$$

Definition: Define the set $\mathrm{Out}(\Delta, \mathrm{M})$ by the formula

$$\mathrm{Out}(\Delta, \mathrm{M}) = (\mathrm{Out}(\mathrm{M})\backslash\mathrm{Hom}_f(\Delta, \mathrm{Out}(\mathrm{M})))/\mathrm{Out}(\Delta).$$

If $\eta \in \mathrm{Hom}_f(\Delta, \mathrm{Out}(\mathrm{M}))$, let $[\eta] = (\mathrm{Out}(\mathrm{M})\eta)\mathrm{Out}(\Delta)$ be the element of $\mathrm{Out}(\Delta, \mathrm{M})$ determined by η.

Let (Γ, N) be a pair such that $[\Gamma, \mathrm{N}] \in \mathrm{Iso}(\Delta, \mathrm{M})$. Let $\mathcal{O} : \Gamma/\mathrm{N} \to \mathrm{Out}_E(\mathrm{N})$ be the homomorphism induced by the action of Γ on N by conjugation. Let $\alpha : \mathrm{N} \to \mathrm{M}$ and $\beta : \Delta \to \Gamma/\mathrm{N}$ be isomorphisms. Then $\alpha_\#\mathcal{O}\beta \in \mathrm{Hom}_f(\Delta, \mathrm{Out}(\mathrm{M}))$.

Let $\alpha' : \mathrm{N} \to \mathrm{M}$ and $\beta' : \Delta \to \Gamma/\mathrm{N}$ are isomorphisms. Observe that

$$\begin{aligned}
\alpha'_\#\mathcal{O}\beta' &= \alpha'_\#\alpha_\#^{-1}\alpha_\#\mathcal{O}\beta\beta^{-1}\beta' \\
&= (\alpha'\alpha^{-1})_\#\alpha_\#\mathcal{O}\beta(\beta^{-1}\beta') \\
&= (\alpha'\alpha^{-1}\mathrm{Inn}(\mathrm{M}))_*(\alpha_\#\mathcal{O}\beta(\beta^{-1}\beta')) \\
&= (\alpha'\alpha^{-1}\mathrm{Inn}(\mathrm{M}))(\alpha_\#\mathcal{O}\beta)(\beta^{-1}\beta').
\end{aligned}$$

Hence $[\alpha_\#\mathcal{O}\beta]$ in $\mathrm{Out}(\Delta, \mathrm{M})$ does not depend on the choice of α and β, and so (Γ, N) determines the element $[\alpha_\#\mathcal{O}\beta]$ of $\mathrm{Out}(\Delta, \mathrm{M})$ independent of the choice of α and β.

Suppose $[\Gamma_i, \mathrm{N}_i] \in \mathrm{Iso}(\Delta, \mathrm{M})$ for $i = 1, 2$, and $\phi : (\Gamma_1, \mathrm{N}_1) \to (\Gamma_2, \mathrm{N}_2)$ is an isomorphism of pairs. Let $\alpha : \mathrm{N}_1 \to \mathrm{N}_2$ be the isomorphism obtained by restricting ϕ, and let $\beta : \Gamma_1/\mathrm{N}_1 \to \Gamma_2/\mathrm{N}_2$ be the isomorphism induced by ϕ. Let $\mathcal{O}_i : \Gamma_i/\mathrm{N}_i \to \mathrm{Out}_E(\mathrm{N}_i)$ be the homomorphism induced by the action of Γ_i on N_i by conjugation for $i = 1, 2$. Then $\mathcal{O}_2 = \alpha_\#\mathcal{O}_1\beta^{-1}$ by Lemma 17. Let $\alpha_1 : \mathrm{N}_1 \to \mathrm{M}$ and $\beta_1 : \Delta \to \Gamma_1/\mathrm{N}_1$ be isomorphisms. Let $\alpha_2 = \alpha_1\alpha^{-1}$ and $\beta_2 = \beta\beta_1$. Then we have

$$(\alpha_2)_\#\mathcal{O}_2\beta_2 = (\alpha_1\alpha^{-1})_\#\alpha_\#\mathcal{O}_1\beta^{-1}\beta_2 = (\alpha_1)_\#\mathcal{O}_1\beta_1.$$

Hence (Γ_1, N_1) and (Γ_2, N_2) determine the same element of $\mathrm{Out}(\Delta, \mathrm{M})$. Therefore there is a function

$$\omega : \mathrm{Iso}(\Delta, \mathrm{M}) \to \mathrm{Out}(\Delta, \mathrm{M})$$

defined by $\omega([\Gamma, \mathrm{N}]) = [\alpha_\#\mathcal{O}\beta]$ for any choice of isomorphisms $\alpha : \mathrm{N} \to \mathrm{M}$ and $\beta : \Delta \to \Gamma/\mathrm{N}$.

Lemma 4. *The set* $\mathrm{Out}(\Delta, \mathrm{M})$ *is finite.*

Proof. The group Out(M) is arithmetic by Theorem 1.1 of [1]. Hence Out(M) has only finitely many conjugacy classes of finite subgroups, cf. §5 of [3]. As Δ is finitely generated there are only finitely many homomorphisms from Δ to a finite group G. Therefore $\mathrm{Out(M)}\backslash\mathrm{Hom}_f(\Delta, \mathrm{Out(M)})$ is finite. Hence $\mathrm{Out}(\Delta, \mathrm{M})$ is finite. □

5. Fiber Cohomology Classes

Consider $\omega : \mathrm{Iso}(\Delta, \mathrm{M}) \to \mathrm{Out}(\Delta, \mathrm{M})$. Suppose $[\Gamma_1, \mathrm{N}_1]$ and $[\Gamma_2, \mathrm{N}_2]$ are in the same fiber of ω. We want to define a class $[\Gamma_1, \mathrm{N}_1; \Gamma_2, \mathrm{N}_2; \alpha, \beta]$ in the cohomology group $H^1(\Gamma_1/\mathrm{N}_1, \mathcal{K}_1)$ where Γ_1/N_1 acts on $\mathcal{K}_1$ by $\mathrm{N}_1(b+B)(v+\overline{I})_\star = (\overline{B}v + \overline{I})_\star$.

Let $\alpha_i : \mathrm{N}_i \to \mathrm{M}$ and $\beta_i : \Delta \to \Gamma_i/\mathrm{N}_i$ be isomorphisms for $i = 1, 2$. Let $\mathcal{O}_i : \Gamma_i/\mathrm{N}_i \to \mathrm{Out}_E(\mathrm{N}_i)$ be the homomorphism induced by the action of Γ_i on N_i by conjugation for $i = 1, 2$. As $\omega([\Gamma_1, \mathrm{N}_1]) = \omega([\Gamma_2, \mathrm{N}_2])$, we have that $[(\alpha_1)_\# \mathcal{O}_1 \beta_1] = [(\alpha_2)_\# \mathcal{O}_2 \beta_2]$. Then there exists α_0 in Aut(M) and β_0 in Aut(Δ) such that $(\alpha_1)_\# \mathcal{O}_1 \beta_1 = (\alpha_0)_\# (\alpha_2)_\# \mathcal{O}_2 \beta_2 \beta_0$. We have that

$$\mathcal{O}_1 = (\alpha_1^{-1} \alpha_0 \alpha_2)_\# \mathcal{O}_2 \beta_2 \beta_0 \beta_1^{-1}.$$

Let $\alpha : \mathrm{N}_1 \to \mathrm{N}_2$ be the isomorphism $\alpha_2^{-1} \alpha_0^{-1} \alpha_1$, and let $\beta : \Gamma_1/\mathrm{N}_1 \to \Gamma_2/\mathrm{N}_2$ be the isomorphism $\beta_2 \beta_0 \beta_1^{-1}$. Then $\mathcal{O}_1 = \alpha_\#^{-1} \mathcal{O}_2 \beta$. Now α induces an isomorphism $\overline{\alpha} : \overline{\mathrm{N}}_1 \to \overline{\mathrm{N}}_2$ defined by $\overline{\alpha}(\overline{\nu}) = \overline{\alpha(\nu)}$ for each ν in N_1. Let $V_i = \mathrm{Span}(\mathrm{N}_i)$ for $i = 1, 2$, and let $\tilde{\alpha} : V_1 \to V_2$ be an affinity such that $\tilde{\alpha} \overline{\mathrm{N}}_1 \tilde{\alpha}^{-1} = \overline{\mathrm{N}}_2$ and $\tilde{\alpha}_* = \overline{\alpha}$, that is, $\tilde{\alpha} \overline{\nu} \tilde{\alpha}^{-1} = \overline{\alpha}(\overline{\nu})$ for each ν in N_1.

Let $\Xi_i : \Gamma_i/\mathrm{N}_i \to \mathrm{Isom}(V_i/\mathrm{N}_i)$ be the homomorphism induced by the action of Γ_i/N_i on V_i/N_i for $i = 1, 2$. Let $\Omega_i : \mathrm{Isom}(V_i/\mathrm{N}_i) \to \mathrm{Out}_E(\mathrm{N}_i)$ be defined so that $\Omega_i(\zeta) = \mathrm{Inn}(\mathrm{N}_i)\hat{\zeta}_*$ where $\hat{\zeta}$ is an isometry of V_i that lifts ζ and $\hat{\zeta}_*$ is the automorphism of N_i defined by $\overline{\hat{\zeta}_*(\nu)} = \hat{\zeta} \overline{\nu} \hat{\zeta}^{-1}$ for $i = 1, 2$. Then we have that $\Omega_i \Xi_i = \mathcal{O}_i$ for $i = 1, 2$. By Lemma 10 of [9], we have that

$$\Omega_1 \Xi_1 = \alpha_\#^{-1} \Omega_2 \Xi_2 \beta = \Omega_1(\tilde{\alpha}_\sharp^{-1} \Xi_2 \beta).$$

Let $\phi : \Gamma_1/\mathrm{N}_1 \to \mathrm{Aff}(V_1/\mathrm{N}_1)$ and $\psi : \Gamma_1/\mathrm{N}_1 \to \mathrm{Aff}(V_1/\mathrm{N}_1)$ be the homomorphisms defined by $\phi = \tilde{\alpha}_\sharp^{-1} \Xi_2 \beta$ and $\psi = \Xi_1$. Then we have that $\phi(g)\psi(g)^{-1}$ is in $\mathcal{K}_1$ for each g in Γ_1/N_1 by Theorem 3 of [9]. As ψ takes values in $\mathrm{Isom}(V_1/\mathrm{N}_1)$ and $\mathcal{K}_1$ is a subgroup of $\mathrm{Isom}(V_1/\mathrm{N}_1)$, we have that ϕ takes values in $\mathrm{Isom}(V_1/\mathrm{N}_1)$.

Let g, h be in Γ_1/N_1, then we have that

$$\begin{aligned}
\phi(gh)\psi(gh)^{-1} &= \phi(g)\phi(h)(\psi(g)\psi(h))^{-1} \\
&= \phi(g)\phi(h)\psi(h)^{-1}\psi(g)^{-1} \\
&= \phi(g)\psi(g)^{-1}\psi(g)\phi(h)\psi(h)^{-1}\psi(g)^{-1} \\
&= (\phi(g)\psi(g)^{-1})\psi(g)_*(\phi(h)\psi(h)^{-1})
\end{aligned}$$

with

$$\psi(\mathrm{N}_1(b+B))_*(v+\overline{I}) = (\overline{b}+\overline{B})_\star (v+\overline{I})_\star (\overline{b}+\overline{B})_\star^{-1} = (\overline{B}v + \overline{I})_\star.$$

Hence the function $\phi\psi^{-1} : \Gamma_1/\mathrm{N}_1 \to \mathcal{K}_1$ is a crossed homomorphism, and so determines a class $[\Gamma_1, \mathrm{N}_1; \Gamma_2, \mathrm{N}_2; \alpha, \beta]$ in $H^1(\Gamma_1/\mathrm{N}_1, \mathcal{K}_1)$, cf. p. 105 of [6].

Let $[\Gamma, \mathrm{N}]$ be a class in $\mathrm{Isom}(\Delta, \mathrm{M})$, and let $V = \mathrm{Span(N)}$. Let $\mathcal{C}$ be the centralizer of $\overline{\mathrm{N}}$ in $\mathrm{Aff}(V)$. By Lemmas 6 and 8 of [9], we have that

$$\mathcal{C} = \{v + \overline{I} : v \in \mathrm{Span}(Z(\mathrm{N}))\}.$$

The group Γ/N acts on $\mathcal{C}$ by $(\mathrm{N}(b+B))(v+\overline{I}) = \overline{B}v+\overline{I}$ and Γ/N acts on $Z(\mathrm{N})$ by $(\mathrm{N}(b+B))(u+I) = Bu+I$. We have a short exact sequence of (Γ/N)-modules

$$0 \to Z(\mathrm{N}) \xrightarrow{\iota} \mathcal{C} \xrightarrow{\kappa} \mathcal{K} \to 0$$

where $\iota(u+I) = u+\overline{I}$ and $\kappa(v+\overline{I}) = (v+\overline{I})_\star$.

Let T be the group of translations of Γ. Then TN/N is a normal subgroup of Γ/N of finite index and TN/N is a subgroup of the group of translations of Γ/N of finite index by Theorem 16 of [8]. The group Γ/N acts on the abelian group TN/N by $(\mathrm{N}(b+B))(a+I) = \mathrm{N}(Ba+I)$, and so TN/N is a (Γ/N)-module. Moreover the group TN/N acts trivially on $\mathcal{C}$.

Lemma 5. *Let $f : \Gamma/\mathrm{N} \to \mathcal{C}$ be a crossed homomorphism, and let $f_{res} : \mathrm{TN}/\mathrm{N} \to \mathcal{C}$ be the restriction of f. Then f_{res} is a homomorphism of (Γ/N)-modules and the class of f in $H^1(\Gamma/\mathrm{N}, \mathcal{C})$ is completely determined by f_{res}.*

Proof. According to [6], p. 354, we have an exact sequence of homomorphisms

$$H^1(\Gamma/\mathrm{TN}, \mathcal{C}) \xrightarrow{inf} H^1(\Gamma/\mathrm{N}, \mathcal{C}) \xrightarrow{res} H^1(\mathrm{TN}/\mathrm{N}, \mathcal{C})^{\Gamma/\mathrm{N}} \to H^2(\Gamma/\mathrm{TN}, \mathcal{C}).$$

The group Γ/TN is finite and $\mathcal{C}$ is a torsion-free, divisible, abelian group, and so $H^i(\Gamma/\mathrm{TN}, \mathcal{C}) = 0$ for $i = 1, 2$ by Corollary IV.5.4 of [6]. Hence

$$res : H^1(\Gamma/\mathrm{N}, \mathcal{C}) \to H^1(\mathrm{TN}/\mathrm{N}, \mathcal{C})^{\Gamma/\mathrm{N}}$$

is an isomorphism. Here $res([f]) = [f_{res}]$. By the Universal Coefficients Theorem (p. 77 of [6]), we have that

$$H^1(\mathrm{TN}/\mathrm{N}, \mathcal{C})^{\Gamma/\mathrm{N}} = \mathrm{Hom}(\mathrm{TN}/\mathrm{N}, \mathcal{C})^{\Gamma/\mathrm{N}}.$$

Here Γ/N acts on $\mathrm{Hom}(\mathrm{TN}/\mathrm{N}, \mathcal{C})$ by $((\mathrm{N}\gamma)h)(x) = (\mathrm{N}\gamma)h(\mathrm{N}\gamma^{-1}x)$ for each $\gamma \in \Gamma$, homomorphism $h : \mathrm{TN}/\mathrm{N} \to \mathcal{C}$, and element $x \in \mathrm{TN}/\mathrm{N}$. Therefore, we have that

$$\mathrm{Hom}(\mathrm{TN}/\mathrm{N}, \mathcal{C})^{\Gamma/\mathrm{N}} = \mathrm{Hom}_{\Gamma/\mathrm{N}}(\mathrm{TN}/\mathrm{N}, \mathcal{C}).$$

Hence $[f_{res}] = \{f_{res}\}$ and f_{res} is a homomorphism of (Γ/N)-modules. Therefore the class of f in $H^1(\Gamma/\mathrm{N}, \mathcal{C})$ is completely determined by f_{res}. □

Lemma 6. *Suppose that $\omega([\Gamma_1, \mathrm{N}_1]) = \omega([\Gamma_2, \mathrm{N}_2])$ with $\mathcal{O}_1 = \alpha_\#^{-1}\mathcal{O}_2\beta$. If the class $[\Gamma_1, \mathrm{N}_1; \Gamma_2, \mathrm{N}_2; \alpha, \beta]$ is in the image of $\kappa_* : H^1(\Gamma_1/\mathrm{N}_1, \mathcal{C}_1) \to H^1(\Gamma_1/\mathrm{N}_1, \mathcal{K}_1)$, then $[\Gamma_1, \mathrm{N}_1] = [\Gamma_2, \mathrm{N}_2]$.*

Proof. Suppose that $[\Gamma_1, \mathrm{N}_1; \Gamma_2, \mathrm{N}_2; \alpha, \beta]$ is in the image of $\kappa_* : H^1(\Gamma_1/\mathrm{N}_1, \mathcal{C}_1) \to H^1(\Gamma_1/\mathrm{N}_1, \mathcal{K}_1)$. Then there is a crossed homomorphism $f : \Gamma_1/\mathrm{N}_1 \to \mathcal{C}_1$ such that

$$\kappa_*([f]) = [\Gamma_1, \mathrm{N}_1; \Gamma_2, \mathrm{N}_2; \alpha, \beta].$$

By Lemma 5, the cohomology class $[f]$ is completely determined by the restriction (Γ_1/N_1)-module homomorphism $f_{res} : \mathrm{T}_1\mathrm{N}_1/\mathrm{N}_1 \to C_1$.

To simplify notation, replace Γ_1/N_1 with Γ_1'. Then $\mathrm{T}_1\mathrm{N}_1/\mathrm{N}_1$ corresponds to $T_1' = \{b'+I' : b+I \in T_1\}$. Moreover Γ_1/N_1 acts on T_1' by $(\mathrm{N}_1(b+B))(b'+I') = B'b'+I'$ and f_{res} corresponds to a homomorphism of (Γ_1/N_1)-modules $f'_{res} : T_1' \to \mathcal{C}_1$.

The group $\mathrm{T}_1\mathrm{N}_1/\mathrm{N}_1$ has finite index in the group of translations of Γ_1/N_1, and so T_1' has finite index in the group of translations of Γ_1'. Hence $\{b' : b+I \in T_1\}$ contains a basis of the vector space $V_1^\perp$. Therefore $f'_{res} : T_1' \to C_1$ induces a linear transformation $L : V_1^\perp \to \mathrm{Span}(Z(\mathrm{N}_1))$ such that $f'_{res}(b'+I') = L(b') + \overline{I}$ for each $b+I$ in T_1 and if $b+B$ is in Γ_1, then $LB' = \overline{B}L$.

Consider the function $h : \Gamma_1/\mathrm{N}_1 \to \mathcal{C}_1$ defined by $h(\mathrm{N}_1(b+B)) = L(b') + \overline{I}$. Then h is a crossed homomorphism. If $b + I$ is in T_1, then

$$h_{res}(\mathrm{N}_1(b+I)) = L(b') + \overline{I} = f'_{res}(b' + I') = f_{res}(\mathrm{N}_1(b+I)).$$

Hence $h_{res} = f_{res}$. Therefore $[h] = [f]$ in $H^1(\Gamma_1/\mathrm{N}_1, \mathcal{C}_1)$ by Lemma 5.

Let $\tilde{\alpha} : V_1 \to V_2$ be the affinity defined above and write $\tilde{\alpha} = \overline{c} + \overline{C}$ with $\overline{c} \in V_2$ and $\overline{C} : V_1 \to V_2$ a linear isomorphism. Define a linear transformation $D : V_1^{\perp} \to \mathrm{Span}(\mathrm{N}_2)$ by $D = \overline{C}L$. If $b + B \in \Gamma_1$, then $DB' = \overline{CBC}^{-1}D$. Let $p_1 : \Gamma_1/\mathrm{N}_1 \to V_1^{\perp}$ be the crossed homomorphism defined by $p_1(\mathrm{N}_1(b+B)) = b'$. Then $h(\mathrm{N}_1\gamma) = \overline{C}^{-1}Dp_1(\mathrm{N}_1\gamma) + \overline{I}$. Observe that $\kappa_*([h]) = [h_\star]$ where $h_\star$ is defined by $h_\star(\mathrm{N}_1\gamma) = (h(\mathrm{N}_1\gamma))_\star$. Thus $h_\star = (\overline{C}^{-1}Dp_1)_\star$ as defined in Theorem 3(3).

Let v be in $\mathrm{Span}(Z(\mathrm{N}_1))$, and let $f_v : \Gamma_1/\mathrm{N}_1 \to \mathcal{K}_1$ be the principal crossed homomorphism determined by $(v+I)_\star$. Then we have that

$$\begin{aligned} f_v(\mathrm{N}_1(b+B)) &= (\mathrm{N}_1(b+B))(v+\overline{I})_\star(v+\overline{I})_\star^{-1} \\ &= (\overline{B}v + \overline{I})_\star(-v+\overline{I})_\star \quad = \quad (\overline{B}v - v + \overline{I})_\star. \end{aligned}$$

Now we have that $[h_\star] = [\Gamma_1, \mathrm{N}_1; \Gamma_2, \mathrm{N}_2; \alpha, \beta]$ in $H^1(\Gamma_1/\mathrm{N}_1, \mathcal{K}_1)$. Hence there exists v in $\mathrm{Span}(Z(\mathrm{N}_1))$ such that

$$(\tilde{\alpha}_\sharp^{-1}\Xi_2\beta)\Xi_1^{-1}f_v = (\overline{C}^{-1}Dp_1)_\star.$$

Let $\tilde{\alpha}_v : V_1 \to V_2$ be the affinity defined by $\tilde{\alpha}_v = \tilde{\alpha}(v+\overline{I})$. Then $\tilde{\alpha}_v\overline{N}_1\tilde{\alpha}_v^{-1} = \overline{N}_2$ and $(\tilde{\alpha}_v)_* = \tilde{\alpha}_*$ by Lemma 6 of [9], and so $(\tilde{\alpha}_v)_* = \overline{\alpha}$. Observe that

$$\begin{aligned} &\big((\tilde{\alpha}_\sharp^{-1}\Xi_2\beta)\Xi_1^{-1}f_v\big)(\mathrm{N}_1(b+B)) \\ &= (\tilde{\alpha}_\sharp^{-1}\Xi_2\beta)(\mathrm{N}_1(b+B))\Xi_1^{-1}(\mathrm{N}_1(b+B))f_v(\mathrm{N}_1(b+B)) \\ &= \tilde{\alpha}_\star^{-1}(\Xi_2\beta)(\mathrm{N}_1(b+B))\tilde{\alpha}_\star(\overline{b}+\overline{B})_\star^{-1}(\overline{B}v - v + \overline{I})_\star \\ &= \tilde{\alpha}_\star^{-1}(\Xi_2\beta)(\mathrm{N}_1(b+B))\tilde{\alpha}_\star(\overline{b}+\overline{B})_\star^{-1}(v - \overline{B}v + \overline{I})_\star^{-1} \\ &= \tilde{\alpha}_\star^{-1}(\Xi_2\beta)(\mathrm{N}_1(b+B))\tilde{\alpha}_\star(v - \overline{B}v + \overline{b} + \overline{B})_\star^{-1} \\ &= \tilde{\alpha}_\star^{-1}(\Xi_2\beta)(\mathrm{N}_1(b+B))\tilde{\alpha}_\star((v+\overline{I})(\overline{b}+\overline{B})(-v+I))_\star^{-1} \\ &= \tilde{\alpha}_\star^{-1}(\Xi_2\beta)(\mathrm{N}_1(b+B))\tilde{\alpha}_\star(v+\overline{I})_\star(\overline{b}+\overline{B})_\star^{-1}(-v+I)_\star \\ &= (-v+I)_\star\tilde{\alpha}_\star^{-1}(\Xi_2\beta)(\mathrm{N}_1(b+B))\tilde{\alpha}_\star(v+\overline{I})_\star(\overline{b}+\overline{B})_\star^{-1} \\ &= (\tilde{\alpha}_v)_\star^{-1}(\Xi_2\beta)(\mathrm{N}_1(b+B))(\tilde{\alpha}_v)_\star(\overline{b}+\overline{B})_\star^{-1} \\ &= \big(((\tilde{\alpha}_v)_\sharp^{-1}\Xi_2\beta)\Xi_1^{-1}\big)(\mathrm{N}_1(b+B)) \end{aligned}$$

Hence we have

$$(\tilde{\alpha}_\sharp^{-1}\Xi_2\beta)\Xi_1^{-1}f_v = ((\tilde{\alpha}_v)_\sharp^{-1}\Xi_2\beta)\Xi_1^{-1}.$$

Thus we have that

$$(\tilde{\alpha}_v)_\sharp^{-1}\Xi_2\beta = (\overline{C}^{-1}Dp_1)_\star\Xi_1.$$

Let $\mathrm{P}_i : \Gamma_i/\mathrm{N}_i \to \Gamma'_i$ be the isomorphism defined by $\mathrm{P}_i(\mathrm{N}_i\gamma) = \gamma'$ for each $i = 1, 2$. Let $\beta' : \Gamma'_1 \to \Gamma'_2$ be the isomorphism so that $\mathrm{P}_2^{-1}\beta'\mathrm{P}_1 = \beta$. Let $\tilde{\beta} : V_1^{\perp} \to V_2^{\perp}$ be an affinity such that $\tilde{\beta}\Gamma'_1\tilde{\beta}^{-1} = \Gamma'_2$ and $\tilde{\beta}_* = \beta'$, that is, and $\tilde{\beta}\gamma'\tilde{\beta}^{-1} = \beta'(\gamma')$ for each γ in Γ_1. Then we have that

$$(\tilde{\alpha}_v)_\sharp^{-1}\Xi_2\mathrm{P}_2^{-1}\tilde{\beta}_*\mathrm{P}_1 = (\overline{C}^{-1}Dp_1)_\star\Xi_1.$$

Therefore there exists an affinity $\phi = c + C$ of E^n such that $\phi(\Gamma_1, \mathrm{N}_1)\phi^{-1} = (\Gamma_2, \mathrm{N}_2)$ with $\overline{\phi} = \tilde{\alpha}_v$, $\phi' = \tilde{\beta}$, and $\overline{C'} = D$ by Theorem 3. Thus $[\Gamma_1, \mathrm{N}_1] = [\Gamma_2, \mathrm{N}_2]$. □

Lemma 7. *Suppose that* $\omega([\Gamma,\mathrm{N}]) = \omega([\Gamma_1,\mathrm{N}_1])$ *with* $\mathcal{O} = (\alpha_1)_\#^{-1}\mathcal{O}_1\beta_1$ *and that* $\omega([\Gamma,\mathrm{N}]) = \omega([\Gamma_2,\mathrm{N}_2])$ *with* $\mathcal{O} = (\alpha_2)_\#^{-1}\mathcal{O}_2\beta_2$. *If* $[\Gamma,\mathrm{N};\Gamma_i,\mathrm{N}_i;\alpha_i,\beta_i]$ *for* $i = 1,2$ *are in the same coset of the image of* $\kappa_* : H^1(\Gamma/\mathrm{N},\mathcal{C}) \to H^1(\Gamma/\mathrm{N},\mathcal{K})$ *in* $H^1(\Gamma/\mathrm{N},\mathcal{K})$, *then* $[\Gamma_1,\mathrm{N}_1] = [\Gamma_2,\mathrm{N}_2]$.

Proof. Let $b + B \in \Gamma$, and let $b_1 + B_1$ be an element of Γ_1 such that

$$\mathrm{N}_1(b_1 + B_1) = \beta_1(\mathrm{N}(b + B)).$$

As $\mathcal{O} = (\alpha_1)_\#^{-1}\mathcal{O}_1\beta_1$, we have that

$$\begin{aligned}
(b+B)_*\mathrm{Inn}(\mathrm{N}) &= (\alpha_1)_\#^{-1}\mathcal{O}_1\beta_1(\mathrm{N}(b+B)) \\
&= (\alpha_1)_\#^{-1}\mathcal{O}_1(\mathrm{N}_1(b_1+B_1)) \\
&= (\alpha_1)_\#^{-1}((b_1+B_1)_*\mathrm{Inn}(\mathrm{N}_1)) \\
&= \alpha_1^{-1}(b_1+B_1)_*\alpha_1\mathrm{Inn}(\mathrm{N}) \\
&= (\tilde{\alpha}_1)_*^{-1}(b_1+B_1)_*(\tilde{\alpha}_1)_*\mathrm{Inn}(\mathrm{N}) \\
&= (\overline{c}_1+\overline{C}_1)_*^{-1}(b_1+B_1)_*(\overline{c}_1+\overline{C}_1)_*\mathrm{Inn}(\mathrm{N}).
\end{aligned}$$

The action of Γ/N on $\mathcal{C}$ is given by $\mathrm{N}(b+B)(u+\overline{I}) = \overline{B}u + \overline{I}$ and is determined by the action of Γ/N on $Z(\mathrm{N})$ induced by conjugation. Now $\mathrm{Inn}(\mathrm{N})$ acts trivially on $Z(\mathrm{N})$, and so the last computation implies that $\overline{B} = \overline{C}_1^{-1}\overline{B}_1\overline{C}_1$ on $\mathrm{Span}(Z(\mathrm{N}))$. Hence the pair of isomorphisms

$$\pi = (\beta_1 : \Gamma/\mathrm{N} \to \Gamma_1/\mathrm{N}_1, (\tilde{\alpha}_1)_*^{-1} : \mathcal{C}_1 \to \mathcal{C})$$

is a change of groups isomorphism in the sense of [6] p. 108. Therefore we have an isomorphism $\pi^* : H^1(\Gamma_1/\mathrm{N}_1,\mathcal{C}_1) \to H^1(\Gamma/\mathrm{N},\mathcal{C})$ defined so that if $f_1 : \Gamma_1/\mathrm{N}_1 \to \mathcal{C}_1$ is a crossed homomorphism, then $\pi^*[f_1] = [\pi^* f_1]$ where $\pi^* f_1 : \Gamma/\mathrm{N} \to \mathcal{C}$ is the crossed homomorphism defined by $\pi^* f_1(x) = (\tilde{\alpha}_1)_*^{-1}(f_1(\beta_1(x)))$.

Likewise the pair of isomorphisms

$$\varpi = (\beta_1 : \Gamma/\mathrm{N} \to \Gamma_1/\mathrm{N}_1, (\tilde{\alpha}_1)_\sharp^{-1} : \mathcal{K}_1 \to \mathcal{K})$$

is a change of groups isomorphism which induces an isomorphism ϖ^* such that the following diagram commutes

$$\begin{array}{ccc}
H^1(\Gamma_1/\mathrm{N}_1,\mathcal{C}_1) & \xrightarrow{\pi^*} & H^1(\Gamma/\mathrm{N},\mathcal{C}) \\
\downarrow (\kappa_1)_* & & \downarrow \kappa_* \\
H^1(\Gamma_1/\mathrm{N}_1,\mathcal{K}_1) & \xrightarrow{\varpi^*} & H^1(\Gamma/\mathrm{N},\mathcal{K}).
\end{array}$$

Now we have that $\omega([\Gamma_1,\mathrm{N}_1]) = \omega([\Gamma_2,\mathrm{N}_2])$ with $\mathcal{O}_1 = (\alpha_2\alpha_1^{-1})_\#\mathcal{O}_2\beta_2\beta_1^{-1}$. Moreover, we have that

$$[\Gamma_1,\mathrm{N}_1;\Gamma_2,\mathrm{N}_2;\alpha_2\alpha_1^{-1},\beta_2\beta_1^{-1}] = [((\tilde{\alpha}_2\tilde{\alpha}_1^{-1})_\sharp^{-1}\Xi_2\beta_2\beta_1^{-1})\Xi_1^{-1}].$$

For $i = 1,2$, we have that

$$[\Gamma,\mathrm{N};\Gamma_i,\mathrm{N}_i;\alpha_i,\beta_i] = [((\tilde{\alpha}_i)_\sharp^{-1}\Xi_i\beta_i)\Xi^{-1}].$$

Let $\gamma \in \Gamma$. Observe that

$$\begin{aligned}
&\big(((\tilde{\alpha}_2)_\sharp^{-1}\Xi_2\beta_2)\Xi^{-1}\big)\big(((\tilde{\alpha}_1)_\sharp^{-1}\Xi_1\beta_1)\Xi^{-1}\big)^{-1}(\mathrm{N}\gamma)\\
&= ((\tilde{\alpha}_2)_\sharp^{-1}\Xi_2\beta_2)(\mathrm{N}\gamma)\Xi(\mathrm{N}\gamma)^{-1}\Xi(\mathrm{N}\gamma)\big(((\tilde{\alpha}_1)_\sharp^{-1}\Xi_1\beta_1)(\mathrm{N}\gamma)\big)^{-1}\\
&= ((\tilde{\alpha}_2)_\sharp^{-1}\Xi_2\beta_2)(\mathrm{N}\gamma)(\tilde{\alpha}_1)_\sharp^{-1}\big(\Xi_1\beta_1(\mathrm{N}\gamma)\big)^{-1}\\
&= ((\tilde{\alpha}_2)_\star^{-1}\Xi_2\beta_2)(\mathrm{N}\gamma)(\tilde{\alpha}_2)_\star(\tilde{\alpha}_1)_\star^{-1}\big(\Xi_1\beta_1(\mathrm{N}\gamma)\big)^{-1}(\tilde{\alpha}_1)_\star\\
&= (\tilde{\alpha}_1)_\sharp^{-1}((\tilde{\alpha}_2\tilde{\alpha}_1^{-1})_\sharp^{-1}\Xi_2\beta_2)(\mathrm{N}\gamma)\big(\Xi_1\beta_1(\mathrm{N}\gamma)\big)^{-1}\\
&= (\tilde{\alpha}_1)_\sharp^{-1}((\tilde{\alpha}_2\tilde{\alpha}_1^{-1})_\sharp^{-1}\Xi_2\beta_2\beta_1^{-1}(\beta_1(\mathrm{N}\gamma))\big(\Xi_1\beta_1(\mathrm{N}\gamma)\big)^{-1}\\
&= \varpi^*\big(((\tilde{\alpha}_2\tilde{\alpha}_1^{-1})_\sharp^{-1}\Xi_2\beta_2\beta_1^{-1})\Xi_1^{-1}\big)(\mathrm{N}\gamma).
\end{aligned}$$

Hence we have that

$$\varpi^*([\Gamma_1,\mathrm{N}_1;\Gamma_2,\mathrm{N}_2;\alpha_2\alpha_1^{-1},\beta_2\beta_1^{-1}]) = [\Gamma,\mathrm{N};\Gamma_2,\mathrm{N}_2;\alpha_2,\beta_2][\Gamma,\mathrm{N};\Gamma_1,\mathrm{N}_1;\alpha_1,\beta_1]^{-1}.$$

The right-hand side of the above equation is in the image of $\kappa_* : H^1(\Gamma/\mathrm{N},\mathcal{C}) \to H^1(\Gamma/\mathrm{N},\mathcal{K})$. Therefore $[\Gamma_1,\mathrm{N}_1;\Gamma_2,\mathrm{N}_2;\alpha_2\alpha_1^{-1},\beta_2\beta_1^{-1}]$ is in the image of $(\kappa_1)_* : H^1(\Gamma_1/\mathrm{N}_1,\mathcal{C}_1) \to H^1(\Gamma_1/\mathrm{N}_1,\mathcal{K}_1)$. Hence $[\Gamma_1,\mathrm{N}_1] = [\Gamma_2,\mathrm{N}_2]$ by Lemma 6. □

6. The relative Bieberbach Theorem

Let $[\Gamma,\mathrm{N}]$ be a class in $\mathrm{Isom}(\Delta,\mathrm{M})$, and let T be the group of translations of Γ. Then Γ/TN is a finite group, since Γ/T is finite. The group TN/N acts trivially on $Z(\mathrm{N})$, TN/N, $\mathcal{C}$, $\mathcal{K}$, and so the action of Γ/N on $Z(\mathrm{N})$, TN/N, $\mathcal{C}$, $\mathcal{K}$ induces an action of Γ/TN on $Z(\mathrm{N})$, TN/N, $\mathcal{C}$, $\mathcal{K}$ making $Z(\mathrm{N})$, TN/N, $\mathcal{C}$, $\mathcal{K}$ into (Γ/TN)-modules.

Lemma 8. *The group $H^1(\Gamma/\mathrm{TN},\mathcal{K})$ is finite.*

Proof. The short exact sequence $0 \to Z(\mathrm{N}) \to \mathcal{C} \to \mathcal{K} \to 0$ of (Γ/TN)-modules induces an exact sequence of cohomology groups

$$H^1(\Gamma/\mathrm{TN},\mathcal{C}) \to H^1(\Gamma/\mathrm{TN},\mathcal{K}) \to H^2(\Gamma/\mathrm{TN},Z(\mathrm{N})) \to H^2(\Gamma/\mathrm{TN},\mathcal{C}).$$

As explained in the proof of Lemma 5, the outside groups are trivial, and so $H^1(\Gamma/\mathrm{TN},\mathcal{K})$ is isomorphic to $H^2(\Gamma/\mathrm{TN},Z(\mathrm{N}))$. The group $H^2(\Gamma/\mathrm{TN},Z(\mathrm{N}))$ is a torsion group by Proposition IV.5.3 of [6]. As $Z(\mathrm{N})$ is a free abelian group of finite rank, the group $H^2(\Gamma/\mathrm{TN},Z(\mathrm{N}))$ is finitely generated. Hence $H^2(\Gamma/\mathrm{TN},Z(\mathrm{N}))$ is finite, and so $H^1(\Gamma/\mathrm{TN},\mathcal{K})$ is finite. □

Lemma 9. *The cokernel of $\kappa_* : H^1(\mathrm{TN}/\mathrm{N},\mathcal{C})^{\Gamma/\mathrm{TN}} \to H^1(\mathrm{TN}/\mathrm{N},\mathcal{K})^{\Gamma/\mathrm{TN}}$ is finite.*

Proof. By the Universal Coefficients Theorem, we have that

$$H^1(\mathrm{TN}/\mathrm{N},\mathcal{C})^{\Gamma/\mathrm{TN}} = \mathrm{Hom}(\mathrm{TN}/\mathrm{N},\mathcal{C})^{\Gamma/\mathrm{TN}} = \mathrm{Hom}_{\Gamma/\mathrm{TN}}(\mathrm{TN}/\mathrm{N},\mathcal{C}),$$

$$H^1(\mathrm{TN}/\mathrm{N},\mathcal{K})^{\Gamma/\mathrm{TN}} = \mathrm{Hom}(\mathrm{TN}/\mathrm{N},\mathcal{K})^{\Gamma/\mathrm{TN}} = \mathrm{Hom}_{\Gamma/\mathrm{TN}}(\mathrm{TN}/\mathrm{N},\mathcal{K}).$$

The short exact sequence $0 \to Z(\mathrm{N}) \to \mathcal{C} \to \mathcal{K} \to 0$ induces an exact sequence

$$0 \to \mathrm{Hom}(\mathrm{TN}/\mathrm{N},Z(\mathrm{N})) \to \mathrm{Hom}(\mathrm{TN}/\mathrm{N},\mathcal{C}) \to \mathrm{Hom}(\mathrm{TN}/\mathrm{N},\mathcal{K}) \to \mathrm{Ext}(\mathrm{TN}/\mathrm{N},Z(\mathrm{N}))$$

by Theorem III.3.4 of [6] (with $R = \mathbb{Z}$). We have that $\mathrm{Ext}(\mathrm{TN}/\mathrm{N},Z(\mathrm{N})) = 0$ by Theorems I.6.3 and III.3.5 of [6], since TN/N is a free abelian group. Hence we have a short exact sequence of (Γ/TN)-modules

$$0 \to \mathrm{Hom}(\mathrm{TN}/\mathrm{N},Z(\mathrm{N})) \to \mathrm{Hom}(\mathrm{TN}/\mathrm{N},\mathcal{C}) \to \mathrm{Hom}(\mathrm{TN}/\mathrm{N},\mathcal{K}) \to 0.$$

Hence we have an exact sequence of cohomology groups

$$H^0(\Gamma/\mathrm{TN}, \mathrm{Hom}(\mathrm{TN}/\mathrm{N}, \mathcal{C})) \to H^0(\Gamma/\mathrm{TN}, \mathrm{Hom}(\mathrm{TN}/\mathrm{N}, \mathcal{K}))$$

$$\to H^1(\Gamma/\mathrm{TN}, \mathrm{Hom}(\mathrm{TN}/\mathrm{N}, Z(\mathrm{N}))),$$

which is equivalent to an exact sequence

$$\mathrm{Hom}_{\Gamma/\mathrm{TN}}(\mathrm{TN}/\mathrm{N}, \mathcal{C}) \to \mathrm{Hom}_{\Gamma/\mathrm{TN}}(\mathrm{TN}/\mathrm{N}, \mathcal{K}) \to H^1(\Gamma/\mathrm{TN}, \mathrm{Hom}(\mathrm{TN}/\mathrm{N}, Z(\mathrm{N}))).$$

The group $H^1(\Gamma/\mathrm{TN}, \mathrm{Hom}(\mathrm{TN}/\mathrm{N}, Z(\mathrm{N})))$ is finite, since $\mathrm{Hom}(\mathrm{TN}/\mathrm{N}, Z(\mathrm{N}))$ is a free abelian group of finite rank. Hence the cokernel of $\mathrm{Hom}_{\Gamma/\mathrm{TN}}(\mathrm{TN}/\mathrm{N}, \mathcal{C})) \to \mathrm{Hom}_{\Gamma/\mathrm{TN}}(\mathrm{TN}/\mathrm{N}, \mathcal{K}))$ is finite. Therefore the cokernel of $\kappa_* : H^1(\mathrm{TN}/\mathrm{N}, \mathcal{C})^{\Gamma/\mathrm{TN}} \to H^1(\mathrm{TN}/\mathrm{N}, \mathcal{K})^{\Gamma/\mathrm{TN}}$ is finite. □

Lemma 10. *The cokernel of* $\kappa_* : H^1(\Gamma/\mathrm{N}, \mathcal{C}) \to H^1(\Gamma/\mathrm{N}, \mathcal{K})$ *is finite.*

Proof. We have a short exact sequence $1 \to \mathrm{TN}/\mathrm{N} \to \Gamma/\mathrm{N} \to \Gamma/\mathrm{TN} \to 1$, and so we have a commutative diagram with horizontal exact sequences (cf. p. 354 of [6])

$$\begin{array}{ccccccc} 0 = H^1(\Gamma/\mathrm{TN}, \mathcal{C}) & \to & H^1(\Gamma/\mathrm{N}, \mathcal{C}) & \to & H^1(\mathrm{TN}/\mathrm{N}, \mathcal{C})^{\Gamma/\mathrm{TN}} & \to & H^2(\Gamma/\mathrm{TN}, \mathcal{C}) = 0 \\ \downarrow \alpha & & \downarrow \beta & & \downarrow \gamma & & \downarrow \\ 0 \to H^1(\Gamma/\mathrm{TN}, \mathcal{K}) & \to & H^1(\Gamma/\mathrm{N}, \mathcal{K}) & \to & H^1(\mathrm{TN}/\mathrm{N}, \mathcal{K})^{\Gamma/\mathrm{TN}} & \to & H^2(\Gamma/\mathrm{TN}, \mathcal{K}) \end{array}$$

where the homomorphism α, β, γ are induced by $\kappa : \mathcal{C} \to \mathcal{K}$. By the snake lemma (Lemma III.5.1 of [5]), we have an exact sequence

$$\mathrm{coker}(\alpha) \to \mathrm{coker}(\beta) \to \mathrm{coker}(\gamma).$$

Now $\mathrm{coker}(\alpha) = H^1(\Gamma/\mathrm{TN}, \mathcal{K})$ is finite by Lemma 8, and $\mathrm{coker}(\gamma)$ is finite by Lemma 9. Hence $\mathrm{coker}(\beta)$ is finite. Thus the cokernel of $\kappa_* : H^1(\Gamma/\mathrm{N}, \mathcal{C}) \to H^1(\Gamma/\mathrm{N}, \mathcal{K})$ is finite. □

Theorem 4. *For each dimension* n*, there are only finitely many isomorphism classes of pairs of groups* (Γ, N) *such that* Γ *is an* n*-space group and* N *is a normal subgroup of* Γ *such that* Γ/N *is a space group.*

Proof. Let m be a positive integer less than n. Let M be an m-space group and let Δ be an $(n-m)$-space group. Let $\mathrm{Iso}(\Delta, \mathrm{M})$ be the set of isomorphism classes of pairs (Γ, N) where N is a normal subgroup of an n-space group Γ such that N is isomorphic to M and Γ/N is isomorphic to Δ. As there are only finitely many isomorphism classes of the groups Δ and M by Bieberbach's theorem [2], it suffices to prove that $\mathrm{Iso}(\Delta, \mathrm{M})$ is finite.

In §4, we defined a function $\omega : \mathrm{Iso}(\Delta, \mathrm{M}) \to \mathrm{Out}(\Delta, \mathrm{M})$ with $\mathrm{Out}(\Delta, \mathrm{M})$ finite by Lemma 4. The fibers of ω are finite by Lemmas 7 and 10. Therefore $\mathrm{Iso}(\Delta, \mathrm{M})$ is finite. □

In view of Theorem 10 of [8], Theorem 4 is equivalent to the following theorem.

Theorem 5. *For each dimension* n*, there are only finitely many affine equivalence classes of geometric orbifold fibrations of compact, connected, flat* n*-orbifolds.*

References

[1] O. Baues and F. Grunewald, Automorphism groups of polycyclic-by-finite groups and arithmetic groups, Publ. Math. Inst. Hautes Études Sci. 104 (2006), 213-268.
[2] L. Bieberbach, Über die Bewegungsgruppen der Euklidischen Rähme II, Math. Ann. 72 (1912), 400-412.
[3] A. Borel, Arithmetic properties of linear algebraic groups, Proc. Int. Congress Math., Stockholm (1992), 10-22.
[4] J. H. Conway, O. D. Friedrichs, D. H. Huson, W. P. Thurston, On three-dimensional space groups, *Beiträge Algebra Geom.* **42** (2001), 475-507.
[5] P. J. Hilton and U. Stammbach, *A Course in Homological Algebra*, Graduate Texts in Math., vol. **4**, Springer-Verlag, New York, Heidelberg, and Berlin, 1971.
[6] S. Mac Lane, *Homology*, Springer-Verlag, New York, 1967.
[7] J.G. Ratcliffe, *Foundations of Hyperbolic Manifolds, Second Edition*, Graduate Texts in Math., vol. **149**, Springer-Verlag, Berlin, Heidelberg, and New York, 2006.
[8] J. G. Ratcliffe and S. T. Tschantz, Fibered orbifolds and crystallographic groups, *Algebr. Geom. Topol.* **10** (2010), 1627-1664.
[9] J. G. Ratcliffe and S. T. Tschantz, On the isometry group of a compact flat orbifold, *Geom. Dedicata* 177 (2015), 43-60.
[10] J. G. Ratcliffe and S. T. Tschantz, The Calabi Construction for Compact Flat Orbifolds, Topology Appl. 178 (2014), 87-106.

Department of Mathematics, Vanderbilt University, Nashville, TN 37240

E-mail address: j.g.ratcliffe@vanderbilt.edu